

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 26, 2016, 2





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies

Series historia naturalis, 26, 2016, 2

ISSN 1408-533X

UDK 5

Letnik 26, leto 2016, številka 2

UREDNIŠKI ODBOR/ COMITATO DI REDAZIONE/ BOARD OF EDITORS:	Dunja Bandelj Mavsar, Nicola Bettoso (IT), Christian Capapé (F), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Mitja Kaligarič, Gregor Kovačič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Alenka Malej, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (A), Tom Turk, Elena Varljen Bužan
Glavni urednik/Redattore capo/ Editor in chief:	Darko Darovec
Odgovorni urednik naravoslovja/ Redattore responsabile per le scienze naturali/Natural Science Editor:	Lovrenc Lipej
Urednica/Redattrice/Editor:	Martina Orlando-Bonaca
Lektor/Supervisione/Language editor:	Polona Šergon (sl.), Petra Berlot (angl.)
Prevajalci/Traduttori/Translators:	Martina Orlando-Bonaca (sl./it.)
Oblikovalec/Progetto grafico/ Graphic design:	Dušan Podgornik, Lovrenc Lipej
Prelom/Composizione/Typesetting:	Grafis trade d.o.o.
Tisk/Stampa/Print:	Grafis trade d.o.o.
Izdajatelj/Editore/Published by:	Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria©
Za izdajatelja/Per Editore/ Publisher represented by:	Salvator Žitko
Sedež uredništva/Sede della redazione/ Address of Editorial Board:	Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran SI-6330 Piran /Pirano, Fornace/Fornace 41, tel.: +386 5 671 2900, fax 671 2901; e-mail: annales@mbss.org, internet: www.zdjp.si

Redakcija te številke je bila zaključena 30. 11. 2016.

Sofinancirajo/Supporto finanziario/ Financially supported by:	Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS)
--	---

Annales - series historia naturalis izhaja dvakrat letno.

Naklada/Tiratura/Circulation:	300 izvodov/copie/copies
--------------------------------------	--------------------------

Revija *Annales series historia naturalis* je vključena v naslednje podatkovne baze: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL).

Vsi članki so prosto dostopni na spletni strani:
<http://zdjp.si/en/cat/zaloznistvo/periodika/annales-series-historia-naturalis/>
 All articles are freely available via website:
<http://zdjp.si/en/cat/zaloznistvo/periodika/annales-series-historia-naturalis/>

VSEBINA / INDICE GENERALE / CONTENTS

EDITORIAL

Hakan KABASAKAL

Annales – a journal specialized in shark research
 Annales – specializirana znanstvena revija
 za raziskave o morskih psih 131

FAUNA IN FLORA

FAUNA E FLORA

FAUNA AND FLORA

Claudio BATTELLI

Morphometric characteristics, vertical distribution
 and density of the limpet *Patella caerulea* L.
 in relation to different substrata of the bay
 of Koper (Gulf of Trieste, northern Adriatic)
 Morfometrične značilnosti, vertikalna razširjenost
 in gostota latvice *Patella caerulea* L.
 glede na različne tipe podlag v Koprskem
 zalivu (Tržaški zaliv, severni Jadran) 145

Amelio PEZZETTA

Le Orchidaceae d'Abruzzo: aggiornamento
 sistematico e nuova check-list
 Kukavičevke dežele Abruci: sistemska
 dopolnitev in nov seznam vrst 157

**Sonja ŠKORNIK, Marija MEZNARIČ
& Mitja KALIGARIČ**

Functional composition of mid-stream gravel bar
 vegetation (Middle Drava River, NE Slovenia)
 Funkcionalna sestava vegetacije prodišč
 srednjega toka reke Drave (SV Slovenija) 171

IHTIOLOGIJA

ITTIOLOGIA

ICHTHYOLOGY

**Pero TUTMAN, Adem HAMZIĆ,
Edhem HASKOVIĆ & Jakov DULČIĆ**

Neretva rudd, *Scardinius plotizza*, Heckel & Kner,
 1858 (Cyprinidae), endemic fish species
 of the Adriatic watershed; biological-ecological
 and conservation traits
 Neretvanska vrsta krapovca *Scardinius plotizza*,
 Heckel & Kner, 1858 (Cyprinidae), endemične
 ribje vrste iz Jadranskega povodja; biološko-
 ekološke in naravovarstvene značilnosti 185

Martina ORLANDO-BONACA & Domen TRKOV

Clinitrachus argentatus (Risso, 1810) (Perciformes:
 Clinidae) – a less known fish species
 in Slovenian coastal waters (Adriatic Sea)
Clinitrachus argentatus (Risso, 1810) (Perciformes:
 Clinidae) – manj znana vrsta rib v slovenskih
 obalnih vodah (Jadransko morje) 191

Murat BILECENOGLU & M. Baki YOKES

Scuba observations reveal a wider distribution
 range for *Thorogobius macrolepis* (Teleostei:
 Gobiidae)
 Potapljaška vzorčevanja odkrivajo širše območje
 razširjenosti veleluskatega glavača
Thorogobius macrolepis (Teleostei: Gobiidae) 197

**Malek ALI, Adib SAAD, Mohammad ALKHATEEB,
Sihem RAFRAFI-NOUIRA & Christian CAPAPÉ**

First record of hollowsnout grenadier *Coelorinchus
 caelorhincus* (Osteichthyes: Macrouridae) from
 the Syrian coast (eastern Mediterranean)
 Prvi podatek o pojavljanju grenadirja *Coelorinchus
 caelorhincus* (Osteichthyes: Macrouridae)
 ob sirski obali (vzhodno Sredozemsko morje) 203

RECENTNE SPREMEMBE V SREDOZEMSKI

BIODIVERZITETI

CAMBIAMENTI RECENTI NELLA BIODIVERSITÀ

MEDITERRANEA

RECENT CHANGES IN THE MEDITERRANEAN

BIODIVERSITY

**Nicola BETTOSO, Giovanni COMISSO
& Petar KRUŽIĆ**

First record of the tripletail *Lobotes surinamensis*
 (Pisces: Lobotidae) in the lagoon of Marano
 and Grado (Gulf of Trieste, northern Adriatic Sea)
 Prvi zapis o pojavljanju vrste *Lobotes surinamensis*
 (Pisces: Lobotidae) v maranski in gradeški
 laguni (Tržaški zaliv, severni Jadran) 209

Hakan KABASAKAL

Historical dispersal of the great white shark,
Carcharodon carcharias, and bluefin tuna,
Thunnus thynnus, in Turkish waters: decline of
 a predator in response to the loss of its prey
 Zgodovinski pregled pojavljanja belega morskega
 volka, *Carcharodon carcharias*, in modroplavutega
 tuna, *Thunnus thynnus*, v turških vodah: upad
 plenilca kot odziv na izgubo plena 213

Tevfik CEYHAN & Okan AKYOL

Additional record of *Alectis alexandrina* (Carangidae) from the north-western Levantine Sea (Fethiye, Turkey)
Dodaten zapis o pojavljanju vrste Alectis alexandrina (Carangidae) iz severozahodnega Levantskega morja (Fethiye, Turčija) 221

Martina ORLANDO-BONACA, Ante ŽULJEVIĆ & Boris ANTOLIĆ

Is the Port of Koper an inhospitable environment for the settlement of non-indigenous macrophytes?
Je koprsko pristanišče negostoljubno okolje za naseljevanje tujerodnih makrofitov? 225

MISCELLANEA

Nusret DREŠKOVIĆ & Ranko MIRIĆ

Bathymetric chart of Lake Derane as the basis for defining the hydro-ecological optimum of the Deransko wetlands
Batimetrija Deranskega jezera kot temelj za določanje hidro-ekološkega optimum Deranskega mokrišča 235

Ali ALAŞ, Dilek TÜRKER & Ahmet ÖKTENER

Occurrence of *Parabrachiella insidosa* (Heller, 1865) and *Parabrachiella merlucii* (Bassett-Smith, 1896) (Copepoda; Lernaeopodidae) in European hake in Turkey
Pojavljanje vrst Parabrachiella insidosa (Heller, 1865) in Parabrachiella merlucii (Bassett-Smith, 1896) (Copepoda; Lernaeopodidae) na osličih v Turčiji 245

Mohamed Ali BEN SMIDA, Rafika FEHRI-BEDOUI, Alëš BOLJE & M'hamed EL CAFSI

Proteins, fatty acids and nutritional value in the muscle of nine marine species commonly consumed in Tunisia (central Mediterranean)
Hranilna vrednost in vsebnost proteinov in maščobnih kislin v mišičnini morskih vrst v Tuniziji (osrednje Sredozemsko morje) 253

OCENE IN POROČILA

RECENSIONI E RELAZIONI
 REVIEWS AND REPORTS

Michael Stachowitsch

Book review: Biogenic formations in the Slovenian Sea

Navodila avtorjem

Istruzioni per gli autori

Instruction to authors

Kazalo k slikam na ovitku

Index to images on the cover

OCCURRENCE OF *PARABRACHIELLA INSIDIOSA* (HELLER, 1865)
AND *PARABRACHIELLA MERLUCCII* (BASSETT-SMITH, 1896) (COPEPODA;
LERNAEPODIDAE) IN EUROPEAN HAKE IN TURKEY

Ali ALAŞ

Department of Biology, A.K.Education Faculty, Necmettin Erbakan University, Konya, Turkey

Dilek TÜRKER

Department of Biology, Science Faculty, Balıkesir University, Balıkesir, Turkey

Ahmet ÖKTENER

³Department of Fisheries, Bandırma Sheep Research Institute, Balıkesir, Turkey
e-mail: ahmetoktener@yahoo.com

ABSTRACT

In this study, Parabrachiella insidiosa (Heller, 1865) and Parabrachiella merluccii (Bassett-Smith, 1896) (Copepoda; Lernaepodidae) found on gill rakers of European hake, Merluccius merluccius (Linnaeus, 1758) (Pisces; Merlucciidae), are reported for the first time from the area of Turkey. Also, some morphological characters of these parasitic copepods are given using photographs and drawings.

Key words: *Parabrachiella*, european hake, Turkey, parasitic, copepod

SEGNALAZIONE DI *PARABRACHIELLA INSIDIOSA* (HELLER, 1865)
E *PARABRACHIELLA MERLUCCII* (BASSETT-SMITH, 1896) (COPEPODA;
LERNAEPODIDAE) NEL MERLUZZO IN TURCHIA

SINTESI

Nello studio viene riportata per la prima volta la presenza di Parabrachiella insidiosa (Heller, 1865) e Parabrachiella merluccii (Bassett-Smith, 1896) (Copepoda; Lernaepodidae) sulle branchiospine del merluzzo europeo Merluccius merluccius (Linnaeus, 1758) (Pisces; Merlucciidae) in Turchia. Gli autori riportano inoltre alcuni caratteri morfologici di questi copepodi parassiti, con l'ausilio di fotografie e disegni.

Parole chiave: *Parabrachiella*, merluzzo, Turchia, parassiti, copepodi

INTRODUCTION

The European hake (*Merluccius merluccius*) occurs in the Atlantic coasts of Europe and western North Africa; northward up to Norway and Iceland, southward down to Mauritania. It is also found in the Mediterranean Sea and along the southern coast of the Black Sea. The European hake is an economically important species. It is one of the principal target species of trawl fishery in the Mediterranean. *M. merluccius* is an important component of the shelf and upper part of the continental slope fish assemblage in the western Mediterranean Sea (Biagi et al., 2002; Busalacchi et al., 2010). It is usually found between 70 m and 370 m of depth, but may also occur within a wider depth range, from inshore waters (30 m) to 1000 m (Cohen et al., 1990).

Although *P. insidiosa* and *P. merluccii* have been recorded from various localities throughout the world (Thompson & Scott, 1903; Scott & Scott, 1913; Barnard, 1955; Capart, 1959; Kabata, 1963; Kabata, 1979; Kabata, 1992; Kabata & Ho, 1981; Kabata, 1986; Radujkovic & Raibaut, 1989; Raibaut et al., 1998; Benkirane et al., 1999; Boualleg et al., 2010), there have been no records on these parasitic copepods from Turkey before.

This paper presents the first records of *P. insidiosa* and *P. merluccii* in Turkey.

MATERIAL AND METHODS

One hundred and fifty specimens of *Merluccius merluccius* (Linnaeus, 1758) (Pisces; Merlucciidae) were collected by local gears from Turkey's North Aegean Sea and Bandırma Bay in 2014. The collected parasites were fixed in 70% ethanol. Some of the specimens were cleared in lactic acid before the dissection of their appendages. The photos were taken with a Canon EOS 1100D camera connected to a microscope. Measurements were taken in millimetres (mm), with a micrometric programme (Pro-way). The parasites' scientific names, synonyms and hosts were checked with WoRMS (Eds) (2016), Froese & Pauly (2016). Kabata (1979) was consulted for terminology. *P. insidiosa* (MNHN-IU-2013-18738) and *P. merluccii* (MNHN-IU-2013-18737) were deposited in the collections of the Museum National d'Histoire Naturelle (MNHN), Paris, France.

RESULTS

***Parabrachiella insidiosa* (Heller, 1865) (Copepoda; Siphonostomatoida; Lernaepodidae)**

All parasites were firmly attached to the gill rakers. The prevalence and mean intensity of parasite were 9.33% and 1.35, respectively. The total number of parasites was 19. The body length of *P. insidiosa* varies from 8 to 12 mm (Fig.1). Trunk longer than broader, with 2 pairs of posterior processes. Cephalothorax and second



Fig. 1: *Parabrachiella insidiosa*, ♀ (scale bar 2 mm).
Sl. 1: *Parabrachiella insidiosa*, ♀ (merilo 2 mm).

maxilla shorter than the trunk. Maxilliped (Fig. 2e) with a robust corpus, myxa bearing single seta. Subchela short, slender with a big spine near mid-length, a spine next to claw base and spinulation on distal part of inner margin, claw well-delimited, with slender tip and pair of teeth near mid-length. Mandible (Fig. 2a) with dental formula P1, S1, P1, S1, P1, S1, B5.

First maxilla exopod (Fig. 2b) from ventral to lateral armed with two subapical setae; endopod with two digitiform papillae, each bearing apical long and short seta. First antenna (Fig. 2c) indistinctly four-segmented, division between first and second segments not always discernible; distal armature with 1 tubercle and 5 visible elements. Second antenna endopod clearly two-segmented (Fig. 2d), distal segment with three spines, two of equal size, the third longer and broader. Exopod larger than endopod. Second maxillae with tips expanded with indistinct 5 lobate processes (more or less separate) (Fig. 2f).

***Parabrachiella merluccii* (Basset-Smith, 1896) (Copepoda; Siphonostomatoida; Lernaepodidae)**

All parasites were firmly attached to the gill rakers. The prevalence and mean intensity of parasite were




Fig. 2: *Parabradiella insidiosa* (♀): (a) Mandible (0.02 mm); (b) first maxilla (0.02 mm); (c) first antenna (0.05 mm); (d) second antenna (0.04 mm); (e) maxilliped (0.035 mm); (f) bulla.

Sl. 2: *Parabradiella insidiosa* (♀): a) spodnja čeljustnica (0,02 mm); (b) prva zgornja čeljustnica (0,02 mm); (c) prva antena (0,05 mm); (d) druga antena (0,04 mm); (e) čeljustna nožica (0,035 mm); (f) bulla.



Fig. 3: *Parabrachiella merluccii* ♀ (scale bar 2 mm).
Sl. 3: *Parabrachiella merluccii* ♀ (merilo 2 mm).

6.6% and 1, respectively. The total number of parasites was 10.

The body length of *P. merluccii* varies from 4 to 7 mm (Fig. 3). Trunk with 2 pairs of posterior processes. Maxilliped (Fig. 4e, f) with a robust corpus, traces of subdivision near basal end. Myxa with fine small spines and a short seta, barb stout half as long as the claw; near mid-length of inner margin with a secondary tooth, with a pair of teeth on claw some distance above inner margin. Mandible (Fig. 4a) with dental formula P1, S1, P1, S1, P1, S1, B5.

First maxilla (Fig. 4b) long, slender with endopod slightly displaced from ventral to lateral surface, long cylindrical distally with two subequal setae and a short seta, exopod short, narrow with two short setae. First antenna (Fig. 4c) four-segmented with slightly inflated basal part and prominent whip at distal end of second segment, border between first and second indistinct, distal segment with well-developed apical armature comprising one tubercle and 5 setae. Second antenna biramous (Fig. 4d). Exopod more prominent and longer than endopod. Exopod bulbous, covered with robust, conical spinules on rounded tip. Endopod two-segmented, armed apically with a longer and broader seta and two short setae.

Spinulation on the anterior-medial side of endopod. Second maxillae with undivided bulla (Fig. 4g).

DISCUSSION

Raubaut et al. (1998) reviewed the occurrence of 226 parasitic copepod species distributed over 88 genera and 20 families of fishes of the Mediterranean Sea. The seven parasitic copepods related to the European hake were listed as *Chondracanthus merluccii*, *Clavella adunca*, *C. stellata*, *Lernaecera lusci*, *Thysanote impudica*, *P. insidiosa* and *P. merluccii* (Kabata, 1992; Tirard et al., 1996; Raubaut et al., 1998; Gaglio et al., 2011).

P. insidiosa is a highly polymorphic species, specific to the genus *Merluccius*. It is reported from the Mediterranean, the North Atlantic Ocean and the Pacific Ocean. *P. insidiosa* has been reported mainly on the Merluccius genus, such as *Merluccius merluccius* (Radujkovic & Raubaut, 1989; Benmansour & Ben Hassine, 1997; Boualleg et al., 2010), *M. australis* (MacKenzie & Longshaw, 1995), *M. gayi peruanus* (Chero et al., 2014), *M. hubbsi* (Etche-goin & Sardella, 1990), *M. capensis* (Krzeptowski, 1980); *M. paradoxus* (Botha, 1986); *M. bilinearis*, *M. productus* (Kabata & Ho, 1981), *M. gayi gayi* (George-Nascimento, 1996), and less frequently on other fish families, including *Chelidonichthys lucerna*, *Trigla lyra* (Raubaut et al., 1998) and *Dicentrarchus labrax* (Brian, 1906). Its morphological variability has caused several scientists to propose new taxa for the parasite of the South African and Argentinian hake species (*M. capensis*, *M. gayi*). Kabata & Ho (1981) proposed to set the morphology of the posterior extremity as one of the criteria of distinction between the various forms of *P. insidiosa*, the other two being host affiliation and geographical distribution.

P. insidiosa f. insidiosa on *M. merluccius* and other species of this genus from the European and African-Atlantic waters bear well-developed posterior processes, while *P. insidiosa f. lageniformis* (on *M. gayi* from the Atlantic coast of South America) and *P. insidiosa f. pacifica* (on *M. hubbsi* and *M. productus* from the Pacific coasts of South and North America, respectively) are devoid of posterior processes according to the Kabata & Ho (1981) trinomial nomenclature.

The present species were compared with drawings by Kabata (1979) and no major difference was observed. The shape of the trunk was longer than wider, and the morphological features of all the dissected specimens, including structure, size of maxillule exopod and endopod, number of teeth and dental formula of mandible, maxilliped spinulation and location of claw and teeth, size of antenna exopod and endopod, number of segment and seta on antennule, permitted the identification of this copepod as *P. insidiosa*. The second maxillae with 5 lobate processes in our samples are the same as in findings by Kabata (1979) and Barnard (1955).

When the 12 host fish species of the *P. insidiosa* parasite were examined according to family characteristics,




Fig. 4: *Parabradiella merluccii* (♀): (a) Mandible (0.02 mm); (b) first maxilla (0.04 mm); (c) first antenna (0.04 mm); (d) second antenna (0.05 mm); (e) maxilliped (0.035 mm); (f) distal part of maxilliped; (g) bulla (0.8 mm).
Sl. 4: *Parabradiella merluccii* (♀): (a) spodnja čeljustnica (0,02 mm); (b) prva zgornja čeljustnica (0,04 mm); (c) prva antena (0,04 mm); (d) druga antena (0,05 mm); (e) čeljustna nožica (0,035 mm); (f) distalni del čeljustne nožice; (g) bulla (0,8 mm).

75% were found to belong to the Merlucciidae, 17% to the Triglidae, 8% to the Moronidae; in terms of habitat 42% of the fish were bathydemersal, 33% demersal, 17% benthopelagic and 8% pelagic-oceanic; as to their feeding habits, all the 12 fish species were denoted in literature as carnivores.

P. merluccii has been reported from the Mediterranean and the North Atlantic Ocean and only on the Merluccius genus, such as *M. merluccius* (Brian, 1906; Scott & Scott, 1913; Delamare Deboutteville, 1950; Papoutsoglou, 1976; Radujkovic & Raibaut, 1989; Benmansour & Ben Hassine, 1998; Benkirane et al., 1999; Boualleg et al., 2010), *M. polli*, *M. paradoxus* (Nunes-Ruivo, 1956; Capart, 1959), *M. capensis* (Grabda & Soliman, 1975). Kabata (1979) found the Thompson & Scott (1903) record on *Sciaena diacanthus* from the Indian Ocean strange. The morphological characters in

the present species are similar to those of Kabata (1963; 1979). For example, the secondary tooth found at near mid-length inner margin of claw, as indicated by the arrow in figure 4f, resembles Kabata's records.

When the 4 host fish species of the *P. merluccii* parasite were examined according to family characteristics, all of them were denoted in literature as belonging to the Merlucciidae family; in terms of habitat, 75% were defined in literature as bathydemersal and 25% as demersal; and according to their feeding habits, all 4 host species were carnivores.

Ten species of the family Lernaepodidae are reported from marine habitats in Turkey, namely, *Clavellotis fallax*, *C. strumosa*, *C. briani*, *Clavellisa scombri*, *Clavella alata*, *Lernaepoda galei*, *T. impudica*, *Parabrachiella bispinosa*, *P. exigua*, *P. hostilis*. They are reported for the first time from Turkey in this study.

POJAVLJANJE VRST *PARABRACHIELLA INSIDIOSA* (HELLER, 1865) IN *PARABRACHIELLA MERLUCCII* (BASSETT-SMITH, 1896) (COPEPODA; LERNAEPODIDAE) NA OSLIČIH V TURČIJI

Ali ALAŞ

Department of Biology, A.K.Education Faculty, Necmettin Erbakan University, Konya, Turkey

Dilek TÜRKER

Department of Biology, Science Faculty, Balıkesir University, Balıkesir, Turkey

Ahmet ÖKTENER

³Department of Fisheries, Bandırma Sheep Research Institute, Balıkesir, Turkey
e-mail: ahmetoktener@yahoo.com

POVZETEK

Avtorji poročajo o prvem zapisu o dveh vrstah zajedalskih ceponožcev in sicer o *Parabrachiella insidiosa* (Heller, 1865) in *Parabrachiella merluccii* (Bassett-Smith, 1896) (Copepoda; Lernaepodidae), ki so jih našli na škrgah osličev *Merluccius merluccius* (Linnaeus, 1758) (Pisces; Merlucciidae) v Turčiji. Podajajo tudi nekatere morfološke značilnosti teh zajedalskih ceponožcev skupaj z ilustracijami in fotografijami.

Ključne besede: *Parabrachiella*, oslič, Turčija, zajedalci, raki ceponožci

REFERENCES

Barnard, K.H. (1955): South African parasitic copepoda. *Anns. S. Afr. Mus.*, 41, 223-312.

Benkirane, O. F., F. Coste & A. Raibaut (1999): On the morphological variability of the attachment organ of Lernaepodidae (Copepoda: Siphonostomatoida). *Folia Parasitol.*, 46, 67-75.

Benmansour, B. & O. K. Ben Hassine (1997): Premiere Mention en Tunisie de Certains Caligidae et Lernaepodidae (Copepoda) parasites de poissons teleosteens. *Ichthyophysiol. Acta*, 20, 157-175.

Benmansour, B. & O. K. Ben Hassine (1998): Preliminary analysis of parasitic copepod species richness among coastal fishes of Tunisia. *Ital. J. Zool.*, 65, 341-344.

Biagi, F., P. Sartor, G. Ardizzone, P. Belcari, A. Belluscio & F. Serena (2002): Analysis of demersal assemblages off the Tuscany and Latium coasts (north-western Mediterranean). *Sci. Mar.*, 66, 233-242.

Botha, L. (1986): Major endoparasites of the Cape hakes *Merluccius capensis* and *M. paradoxus*, with brief notes on some conspicuous ectoparasites. *S. Afr. J. Marine. Sci.*, 4, 45-49.

- Boualleg, C., M. Seridi, N. Kaquachi, Y. Quiliquini & M. Bensouillah (2010):** Les Copépodes parasites des poissons téléostéens du littoral Est-algérien. Bull. Inst. Sci, Rabat, Sci. Vie, 32, 65-72.
- Brian, A. (1906):** Copepodi parassiti dei Pesci d'Italia. Stab. Tipo-Litografico R. Istituto Sordomuti, Genova, 187 pp.
- Busalacchi, B., P. Rinelli, F. De Domenico, A. Profeta, F. Perdichizzi & T. Bottari (2010):** Analysis of demersal fish assemblages off the Southern Tyrrhenian Sea (central Mediterranean). Hydrobiologia, 654, 111-124.
- Capart, A. (1959):** Copépodes parasites. Résult scient. Expéd. océanogr. Belge Eaux côt. Africaines Atlantique Sud. (1948-1949), 3, 55-126.
- Chero, J., C. Cruces, J. Iannacone, G. Sáez, L. Alvaríño, C. Rodríguez, H. Rodríguez, E. Tuesta, A. Pacheco & N. Huamani (2014):** Parasitological indices of the Peruvian Hake *Merluccius gayi peruanus* Ginsburg, 1954 (Perciformes: Merlucciidae) acquired at the fishing terminal of Ventanilla, Callao, Peru. Neotrop. Helminthol., 8, 141-162.
- Cohen, D. M., T. Inada, T. Iwamoto & N. Scialabba (1990):** Gadiform fishes of the world (Order Gadiformes). An annotated and illustrated Catalogue of Cods, Hakes, Grenadiers and other Gadiform fishes Known to Date. FAO Species catalogue.vol. 10., Synopsis 125, 10-442.
- Delamare-Deboutteville, C. (1950):** Copepodes parasites des poissons de Banyuls. Vie Milieu, 1, 305-309.
- Etchegoin, J. A. & N. H. Sardella (1990):** Some ecological aspects of the copepod parasites of the common hake, *Merluccius hubbsi*, from the Argentine-Uruguayan Common fishing zone. Int. J. Parasitol., 20, 1009-1014.
- Froese, R. & D. Pauly (2016):** FishBase. Available on line at: www.fishbase.org, (Last accessed 11 December 2016).
- Gaglio, G, T. Bottari, P., Rinelli, F. Marino, D. Macri & S. Gianetto (2011):** Prevalence of *Clavella stellata* (Copepoda: Lernaepodidae) (Kroyer 1838) in the European hake (*Merluccius merluccius*) (L. 1758), South Tyrrhenian Sea (Central Mediterranean). J. Appl. Ichthyol., 27, 136-138.
- George-Nascimento, M. (1996):** Populations and assemblages of parasites in hake, *Merluccius gayi*, from the southeastern Pacific Ocean: stock implications. J. Fish Biol., 48, 557-568.
- Grabda, J. & I. A. F. M. Soliman (1975):** Copepods-Parasites of the Genus *Merluccius* from the Atlantic Ocean and Mediterranean Sea. Acta Ichthyol. Piscat., 5, 31-39.
- Kabata, Z. (1963):** The Second Antenna in the Taxonomy of Clavellinae (Copepoda, Lernaepodidae). Crustaceana, 6, 5-14.
- Kabata, Z. (1979):** Parasitic Copepoda of British Fishes. Ray Society Publications, The British Museum, London, 152, 468 pp.
- Kabata, Z. (1986):** Redescriptions of and comments on four little-known Lernaepodidae (Crustacea: Copepoda). Can. J. Zool., 64, 1852-1859.
- Kabata, Z. (1992):** Copepods parasitic on fishes. Synopses of the British Fauna (Edited by Doris M.Kermack, Barnes, R.S.K., Crothers, J.H.) The Linnean Society of London and the Estuarine and Coastal Sciences Association. Universal Book Services/Dr.W. Backhuys, The Netherlands, 264 pp.
- Kabata, Z. & J. S. Ho (1981):** The origin and dispersal of hake (genus *Merluccius*: Pisces: Teleostei) as indicated by its copepod parasites. Oceanogr. Mar. Biol. Annu. Rev., 19, 381-404.
- Krzeptowski, M. (1980):** Occurrence of larval nematode *Anisakis simplex*, larval cestode *Hepatoxylon trichiuri*, and parasitic copepod *Parabrachiella australis* in juvenile *Merluccius capensis* off Namibia. Acta Ichthyol. Piscat., 10, 35-44.
- MacKenzie, K. & M. Longshaw (1995):** Parasites of the hakes *Merluccius australis* and *M. hubbsi* in the waters around the Falkland Islands, southern Chile, and Argentina, with an assessment of their potential value as biological tags. Can. J. Fish. Aquat. Sci., 52 (1), 213-224.
- Nunes-Ruivo, L. (1956):** Copepodes parasitas de peixes dos mares de Angola. Anais Junta Invest. Ultramar, 9 (2), 9-45.
- Papoutsoglou, S. E. (1976):** Metazoan parasites of fishes from Saronicos Gulf, Athens, Greece. Thalassographica, 1(1), 69-91.
- Radujkovic, B. & A. Raibaut (1989):** Parasites des poissons marins des côtes du Monténégro: Copépodes. Acta Adriat., 28, 237-278.
- Raibaut, A., C. Combes & F. Benoit (1998):** Analysis of the parasitic copepod species richness among Mediterranean fish. J. Marine Syst., 15, 185-206.
- Scott, T. & A. Scott (1913):** The British parasitic Copepoda. Ray Society, London, Volumes I and II: 257 pp.
- Thompson, I. C. & A. Scott (1903):** Report on the Copepoda collected by Professor Herdman, at Ceylon, in 1902. In: Herdman, W.A. (ed.). Report to the Government of Ceylon on the Pearl Oyster Fisheries of the Gulf of Manaar, 1, 227-307.
- Tirard, C., F. Thomas, A. Raibaut & F. Renaud (1996):** The distribution and abundance of Lernaecera lusci (Copepoda) on hake (*Merluccius merluccius*) and bib (*Trisopterus luscus*) (Teleostei). Int. J. Parasitol., 26, 1387-1392.
- WoRMS (Eds) (2016):** World Register of Marine Species. Available on line at: <http://www.marinespecies.org> (Last accessed 11 December 2016).