

A Faunistic Study on the Freshwater Copepoda (Crustacea) of Balıkesir

¹Serdar SÖNMEZ*, ¹Serdar SAK, ¹Alp ALPER and ²Süphan KARAYTUĞ

¹ Balıkesir University, Faculty of Arts and Science, Department of Biology, 10145, Balıkesir, Turkey.

² Mersin University, Faculty of Arts and Science, Department of Biology, 33343, Mersin, Turkey.

* Corresponding Author
e-mail: sonmezserdar@gmail.com

Received: March 02, 2008

Accepted: April 29, 2008

Abstract

Freshwater copepods of Balıkesir Province (Turkey) were investigated by sampling 26 various freshwater bodies and a total of 12 species were identified. *Megacyclops viridis*, *Macrocyclus albidus*, *Mesocyclus leuckarti*, *Paracyclops chiltoni*, *Bryocamptus (B.) minutus*, *B. (R.) pygmaeus* and *(B.) tarnogradskyi* were recorded for the first time from Balıkesir. *Bryocamptus (B.) tarnogradskyi* was also a new record for the Turkish inland waters and it was partially described.

Key words: Fauna, copepod, *Bryocamptus (B.) tarnogradskyi*, Taxonomy, Turkey

INTRODUCTION

The sub-class Copepoda contains over 11500 species [1,2] and constitutes a widely distributed group of crustaceans with free-living, parasitic and associated forms [2]. Members of the families Cyclopidae in the Cyclopoida, Canthocamptidae in the Harpacticoida, and the Diaptomidae in the Calanoida are particularly successful in all kinds of freshwater habitats and they play a vital role in the food web as a part of zooplanktonic community [3].

Considerable amount of studies were carried out on zooplankton of Turkish inland waters since early 1900's. Ustaoglu [4] presented a checklist by compiling about 115 papers published between 1940 and 2004. After Ustaoglu's [4] checklist, additional taxonomical and ecological works were published on zooplankton of ponds [5,6], rivers [7], dam lakes [8-12], mountain lakes [13], natural lakes and springs [14-19].

On the other hand, freshwater copepods of Balıkesir have not sufficiently been studied so far. Some copepod records were given only from İkizcetepeler Dam Lake [20] and Manyas Lake [21-24]. Therefore, 26 various freshwater bodies of Balıkesir were sampled in order to contribute to the copepod fauna of Balıkesir and Turkey.

The samples were collected, by using 100 µm mesh sized plankton net, from 26 stations (Fig. 1) within the border of Balıkesir province between 2001 and 2005. The specimens were instantly preserved in 4% formaldehyde and later transferred into alcohol. The specimens were dissected using Olympus SZX12 or SZX16 stereomicroscopes and were identified by using Olympus BX-50 or BX-51 equipped with differential interference contrast microscopes. Fragments of coverslip were placed between the slide and the coverslip to avoid crushing and to facilitate rotation, which allowed the appendages to be viewed from all angles. All drawings were made with the aid of a drawing tube attached to Olympus BX-51 microscope. The descriptive terminology is adopted from Huys et al. [25]. Abbreviations used in the text are: ae, aesthetasc; P1-P6, for swimming legs 1-6; exp (enp)-1 (-2, -3) to denote the proximal

(middle, distal) segment of a ramus. The sampling dates and localities are given in Table 1. The specimens were identified according to Einsle [26], Karaytuğ [27], Ueda and Reid [28], Borutzky [29], Damian-Georgescu [30], Wells [31]. Martin and Davis [32] were followed for higher classification of the species.

MATERIAL and METHODS




Figure 1. Sampling localities; 1. A pond in Kapıdağ Peninsula, 2. Karadere (Manyas), 3. Manyas Lake, 4. Söve pond, 5. Çaylak waterfall (Susurluk), 6. Ilıca pond, 7. Armutalan pond, 8. Şamlı pond, 9. Halkapınar pond, 10. Kavaklı pond, 11. Karakol pond, 12. İbirler pond, 13. Köteyli dam lake, 14. Değirmenboğazi stream, 15. Small creek in Kazdağı (near Kalkım), 16. Temporary pools in Balıkesir fair, 17. Small water pool near Yeni Mahalle (Akçay), 18. Bostancı I stream, 19. Bostancı II stream, 20. Pamukçu (Ilıca) stream, 21. A fountain through in Yeşilköy (near Eski Balath), 22. Simav brook, 23. Yörücekler regulator, 24. Kocabay pond, 25. Çaygören dam lake.

RESULTS

A total of 12 species were determined (Table 1). The species are as follows:

Station No.	Dates	Cyclopoida								Harpacticoida			
		<i>A. robustus</i>	<i>M. viridis</i>	<i>E. serrulatus</i>	<i>C. vicinus</i>	<i>M. albidus</i>	<i>T. prasinus</i>	<i>M. leuckarti</i>	<i>P. chiltoni</i>	<i>B. (B.) minutus</i>	<i>B. (B.) tarnogradskyi</i>	<i>B. (R.) pygmaeus</i>	<i>N. hibernica</i>
1	18.05.2001			+									
2	07.04.2004	+				+							
3	01.05.2004	+	+										
4	09.06.2001	+			+								
5	07.04.2004	+											
6	09.06.2001			+		+	+						
7	09.06.2001	+											
8	04.06.2001	+											
9	09.06.2001	+											
10	09.06.2001	+											
11	09.06.2001	+											
12	09.06.2001	+											
13	09.06.2001			+		+	+	+					
14	09.06.2001	+		+		+							
14	28.07.2001						+	+					
15	12.05.2005	+		+		+	+			+	+	+	
16	09.10.2003	+											
17	17.04.2005		+	+									
18	13.04.2001	+	+										
19	16.06.2002					+							+
20	10.10.2001	+				+							
21	08.11.2002								+				
22	16.06.2002	+	+	+									
23	16.06.2002	+											
24	16.06.2002				+								
25	16.06.2002	+											

Table 1. List of identified species regard to the stations and dates

- Class: Maxillopoda Dahl, 1956
- Subclass: Copepoda Milne-Edwards, 1840
- Infraclass: Neocopepoda Huys & Boxshall, 1991
- Superorder: Podoplea Giesbrecht, 1882
- Order: Cyclopoida Burmeister, 1834
- Family: Cyclopidae Dana, 1846
- Subfamily: Eucyclopinæ Kiefer, 1927

Eucyclops serrulatus (Fischer, 1851)

Material Examined: 12 ♀♀, 3 ♂♂ (St.1), 2 ♀♀ (st6), 1 ♀ (st13), 6 ♀♀, 4 ♂♂ (st14), 1 ♀ (st15), 4 ♀♀, 2 ♂♂ (st17), 20 ♀♀, 20 ♂♂ (st22).

Macrocyclus albidus (Jurine, 1820)

Material Examined: 1 ♀ (st2), 1 ♀ (st6), 5 ♀♀, 1 ♂ st(13), 1 ♀ (st14), 3 ♀♀ (st15), 20 ♀♀, 20 ♂♂ (st19), 20 ♀♀, 20 ♂♂ (st20).

Paracyclops chiltoni (Thomson, 1882)

Material Examined: 13 ♀♀, 2 ♂♂ (st21)

Tropocyclops prasinus (Kiefer, 1978)

Material Examined: 1 ♀ (st6), 4 ♀♀, 2 ♂♂ (st13), 20 ♀♀, 20 ♂♂ (st14), 30 ♀♀, 30 ♂♂ (st15).

Subfamily : Cyclopinae Kiefer, 1927

Acanthocyclops robustus (G.O. Sars, 1863)

Material Examined: 10 ♀♀, 10 ♂♂ (st2), 20 ♀♀, 20 ♂♂ (st3), 20 ♀♀, 20 ♂♂ (st4), 8 ♀♀ (st5), 2 ♀♀, 1 ♂ (st7), 30 ♀♀, 30 ♂♂ (st8), 20 ♀♀, 20 ♂♂ (st9), 11 ♀♀, 4 ♂ (st10), 10 ♀♀, 10 ♂♂ (st11), 2 ♀♀, 10 ♂♂ (st12), 2 ♀♀ (st14), 20 ♀♀, 20 ♂♂ (st15), 3 ♀♀, 8 ♂♂ (st16), 20 ♀♀, 20 ♂♂ (st18), 20 ♀♀, 20 ♂♂ (st20), 20 ♀♀, 20 ♂♂ (st22), 16 ♀♀, 1 ♂ (st23), 20 ♀♀, 20 ♂♂ (st25).

Megacyclops viridis (Jurine, 1820)

Material Examined: 20 ♀♀, 20 ♂♂ (st3), 10 ♀♀, 5 ♂♂ (st17), 20 ♀♀, 20 ♂♂ (st18), 20 ♀♀, 20 ♂♂ (st22).

Mesocyclops leuckarti (Claus, 1857)

Material Examined: 30 ♀♀, 10 ♂♂ (st13), 20 ♀♀, 20 ♂♂ (st14)

Cyclops vicinus Uljanin, 1875

Material Examined: 20 ♀♀, 20 ♂♂ (st4), 20 ♀♀, 20 ♂♂ (st24).

Order: Harpacticoida

Family: Ameiridae Sars, 1911

Nitokra hibernica (Brady, 1880)

Material Examined: 4 ♀♀, 5 ♂♂ (st19).

Family: Canthocamptidae G.O. Sars, 1906

Bryocamptus (Rheocamptus) pygmaeus (G.O. Sars, 1863)

Material Examined: 3 ♀♀, 5 ♂♂ (st15).

Bryocamptus (Bryocamptus) minutus (Claus, 1863)

Material Examined: 1 ♀ (dissected), (st15)

Bryocamptus (B.) tarnogradskyi Borutzky, 1934

Material Examined: 1 ♀ (dissected), (st15).




Figure 2. *Bryocamptus (B.) tarnogradskyi* ♀ A. Habitus, dorsal; B. Anal operculum and caudal ramus, dorsal; C. P1, anterior; D. P3 endopod, anterior; E. P4 endopod, anterior; F. P5, anterior.

Supplementary description of female *B. (B.) tarnogradskyi*: Total body length from tip of rostrum to posterior margin of caudal rami 684 µm. Ornamentation of the body as figured (Fig. 2A). Cephalosome with smooth margin and with medially constricted dorsal hyaline window. Hyaline frills of thoracic somites plane while those of abdominal somites denticulated. Urosomites with rows of spinules posterolaterally. Antennule 8-segmented (Fig. 2A). Exopod of antenna 2-segmented, distal segment with 3 setae. Anal somite (Fig. 2B) with row of spinules on posteroventral margin; anal operculum with 6 well developed spinules. Length/width ratio of caudal rami 1.25 (Fig. 2B). Caudal rami ornamented with spinules; two spinules near the base of seta II, a large spinule near the base of seta III and a group of spinules around inner distal margin as figured (Fig. 2B). Each ramus with seven setae; seta I small; seta VII naked, located dorsally and triarticulated at base. P1-P4 with 3-segmented exopod. P1, P2 and P3 (Fig. 2D) with 3-segmented endopod, P4 with 2-segmented endopod (Fig. 2E). P1 endopod longer than exopod; first endopodal segment slightly exceeding exp-2 (Fig. 2C). Setal formula of swimming legs as follows:

Exopod	Endopod	
P1	0.1.121	1.1.120
P2	0.1.132	1.1.121
P3	0.1.223	1.1.221
P4	0.1.223	1.2.21

P5 (Fig. 2F) outer basal seta long and naked with a spinule at base. Baseoendopod with 5 spinulose setae (innermost seta smallest) and with 2 small setae at outer margin. Exopod about 1.8 times longer than broad; with 1 naked outer seta with a spinule row near base, 1 spinulose seta at outer distal corner and 3 terminal setae (2 naked and 1 spinulose); inner margin with 3 spinules.

Remarks: Although our specimen has an extra seta on the baseoendopod of P5 (arrowed in Fig. 2F) and has 6 spinules on the distal margin of anal operculum, it can best be attributable to *Bryocamptus (B.) tarnogradskyi*. No other detailed comparison can be made with the published descriptions of *B. (B.) tarnogradskyi* [29,33,34] because of the inadequacy in the descriptions. The genus *Bryocamptus* Chappuis, 1929 requires urgent worldwide revision.

DISCUSSION

Examination of the previous literature on the zooplankton of Turkish inland waters revealed that only İkizcetepeler Dam Lake [20] and Manyas Lake [21-24] were previously investigated in Balıkesir province by various authors. The Manyas Lake was studied by Noodt [23], Kiefer [22], Demirhindi [6], Ustaoglu and Balık [9]. Noodt [23] recorded *Nitokra hibernica*; Kiefer [22] reported *Eucyclops serrulatus* and *Mesocyclops leuckarti aequatorialis* (now synonym of *Mesocyclops aequatorialis*); Demirhindi [21] reported *Cyclops agilis* = *C. serrulatus* (now synonyms of *E. serrulatus*), *C. vernalis* (now synonym of *Acanthocyclops robustus*), *C. leuckarti aequatorialis* (now synonym of *Mesocyclops aequatorialis*), *C. vicinus* and *Calanipeda aquaedulcis*; Ustaoglu and Balık [9] determined *C. vicinus*, *A. robustus* and *N. hibernica*. Alper et al. [5] studied the Cladocera and Copepoda fauna of İkizcetepeler Dam Lake and they reported 4 copepod species; *C. vicinus*, *A. robustus*, *Tropocyclops prasinus* and *E. serrulatus*. So, 7 species have been reported from Balıkesir so far according to the previous literature. On the other hand, 12 copepod species were identified in our study, and 7 of which (*Megacyclops viridis*, *Macrocyclus albidus*, *Mesocyclops leuckarti*, *Paracyclops chiltoni*, *B. (B.) minutus*, *B. (R.) pygmaeus* and *B. (B.) tarnogradskyi*) are new records for Balıkesir province.

The recent addition of some new copepod records to the Turkish inland waters have proved that there are still many copepod species waiting to be reported especially from mountain lakes, rivers and from cryptic habitats. The following records of copepods have recently been added to Copepoda fauna of Turkey. Ustaoglu et al. [13] studied the zooplankton of 16 mountain lakes in the Taurus Range and found *Arctodiaptomus (Rh.) alpinus*. Bozkurt [18] reported the harpacticoid copepods, *Phyllognathopus viguieri* (Maupas, 1892) and *Leptocaris brevicornis* (Van Douwe, 1904) from the lakes Gölbaşı and Gökent respectively. Güher and Kırgız [19] found two species of darcythompsoniid harpacticoids, *Horsiella brevicornis* (Van Douwe, 1904) and *Horsiella trisetosa* Kunz, 1935. Güher and Kırgız [19] claimed that these two species were new records for the Turkish fauna. But, Kunz [35] considered *Leptocaris* T. Scott and *Horsiella* Gurney as synonymous due to the discovery of several species constituting transitions between the two genera and hence retained only the genus *Leptocaris*. Therefore these

two species should be cited as *Leptocaris brevicornis* (Van Douwe, 1904) and *Leptocaris trisetosus* (Kunz, 1935).

The specious genus *Bryocamptus* contains 118 species and subspecies worldwide and they were collected from almost all kinds of freshwater bodies [2]. Four species of *Bryocamptus* have previously been reported from Turkish inland waters: *B. (R.) zschokkei* (Schmeil, 1893) [36], *B. (R.) pygmaeus* [4, 36], *B. (R.) thylops* (Mrazek, 1893) [36] and *B. (B.) minutus* [4,5,36]. Among the three *Bryocamptus* species identified in this study, *B. (B.) tarnogradskyi* has not previously been reported from Turkey and thus it is a new record for the Turkish fauna. *B. (B.) tarnogradskyi* was originally described from Teberda, Caucasus [34] and was subsequently reported from Europe [37].

REFERENCES

- [1] Humes A.G. 1994. How many copepods? *Hydrobiologia*. 292-293(1): 1-7.
- [2] Boxshall G., Halsey S.H. 2004. *An Introduction to Copepod Diversity*. The Ray Society, London. 966pp.
- [3] Huys, R., Boxshall G.A. 1991. *Copepod evolution*. The Ray Society. 468pp.
- [4] Ustaoglu M.R. 2004. A Check-list for Zooplankton of Turkish Inland Waters. *E.U. Journal of Fisheries & Aquatic Sciences*. 21: 191-199.
- [5] Bozkurt A., Dural M. 2005. Topboğazi Göleti (Hatay) Zooplanktonunun Vertikal Göçü. *Türk Sucul Yaşam Dergisi*. 4: 104-109.
- [6] Güher H, Erdoğan S. 2008. Alıç Göleti Perifitik Zooplankton (Cladocera, Copepoda, Rotifera) Türleri Üzerine Bir Araştırma. *Journal of Fisheries Sciences.com*. 2(3): 516-523.
- [7] Ziya M., Göksu L., Bozkurt A., Taşdemir M., Sarıhan E. 2005. Asi Nehri (Hatay, Türkiye) Cladocera ve Copepoda (Crustacea) Faunası. *E.U. Journal of Fisheries & Aquatic Sciences* 22(1-2):17- 19.
- [8] Yiğit S., Altındağ A. 2005. A Taxonomical Study on the Zooplankton Fauna of Hirfanlı Dam Lake (Kırşehir), Turkey. *G.U. Journal of Science*. 18(4): 563-567.
- [9] Tellioğlu A., Yılmaztürk Y. 2005. Keban Baraj Gölü, Pertek Bölgesi'nin Kladoser ve Kopepod Faunası Üzerine Taksonomik Bir Çalışma. *E.U. Journal of Fisheries & Aquatic Sciences*. 22 (3-4): 431- 433
- [10] Bekleyen A. 2006. Devegeçidi Baraj Gölü'nün (Diyarbakır) Cladocera ve Copepoda (Crustacea) Faunası. *E.U. Journal of Fisheries & Aquatic Sciences*. 23(3-4): 413-415.
- [11] Kaya M., Altındağ A. 2007. Zooplankton Fauna and Seasonal Changes of Gelingüllü Dam Lake (Yozgat, Turkey) *Turkish Journal of Zoology*. 31: 347-351.
- [12] Bozkurt A. 2008. Birecik Baraj Gölü Zooplanktonunun Vertikal Dağılımı. *Journal of Fisheries Sciences.com*. 2(3): 332-342.
- [13] Ustaoglu M.R., Balık S., Mis D. Ö., Aygen C. 2005. The Zooplankton of Some Mountain Lakes in the Taurus Range (Turkey). 2005. *Zoology in the Middle East*. 34: 101-108
- [14] Aygen C., Balık S. 2005. Işıklı Gölü ve Kaynaklarının (Çivril-Denizli) Crustacea Faunası. *E.U. Journal of Fisheries & Aquatic Sciences*. 22(3-4): 371-375.
- [15] Türkmen M., Naz M., Mete Z. 2006. Dinler Gölbaşı Gölü'nün Zooplankton Tür Kompozisyonu ve Biyomasi (Hatay, Türkiye). *E.U. Journal of Fisheries & Aquatic Sciences*. 23(1/1): 163-167.
- [16] Ustaoglu M.R., Balık S., Aygen C., Mis D. Ö. 2006. Akgöl'ün (Selçuk, İzmir) Kladoser ve Kopepodları. *E.U. Journal of Fisheries & Aquatic Sciences*. 23 (1/1): 169-172.
- [17] Bozkurt A. 2006. Yenişehir Gölü (Reyhanlı, Hatay) Zooplanktonu. *E.U. Journal of Fisheries & Aquatic Sciences*. 23 (1/1): 39-43.
- [18] Bozkurt A. 2007. Two New Records of Harpacticoid Copepods for Turkish Inland Waters (Copepoda, Harpacticoida) *Phyllognathopus viguieri* (Maupas, 1892) and *Leptocaris brevicornis* (Van Douwe, 1904). *Crustaceana*. 80(9): 1033-1042.
- [19] Güher H., Kırgız T. 2007. Gala Gölü Milli Parkı'nda Makrofitler İle Mikrocrustacea (Cladocera, Copepoda) İlişkisi Üzerine Bir Araştırma. *Trakya University Journal of Science*. 8(2): 109-114.
- [20] Alper A., Çelebi E., Çam H., Karaytuğ S. 2007. Cladocera and Copepoda (Crustacea) Fauna of İkizcetepeler Dam Lake (Balıkesir, Turkey). *Turkish Journal of Fisheries and Aquatic Sciences*. 7: 1-73.
- [21] Demirhindi Ü. 1972. The Preliminary Planktonic Investigations in the Coastal Lagoons and Several Brackish Water Lakes of Turkey. *İstanbul Üniversitesi Fen Fakültesi Mecmuası*. 37: 205-232.
- [22] Kiefer F. 1955. Freilebende Ruderfusskrebse (Curstacea Copepoda) aus Türkischen Binnengewassern. II. Cyclopoida und Harpacticoida. *ibid.* B, 2(4): 108-123.
- [23] Noodt W. 1954. Copepoda Harpacticoida aus dem limnischen Mesopsammal der Türkei. *Publ. Inst. Hydrobiol. Univ. İstanbul*. 2(1): 27-40.
- [24] Ustaoglu M.R., Balık S. 1990. Zooplankton of Lake Kuş (Bandırma) (in Turkish). *X. Ulusal Biyoloji Kongresi*, 11-19.
- [25] Huys R., Gee J. M., Moore C. G., Hamond R. 1996. Marine and brackish water harpacticoid copepods. Part 1. *In: Barnes, R.S.K. & J.H. Crothers (Eds), Synopses of the British Fauna (New Series)*, 51: i-viii, 1-352. Field Studies Council, Shrewsbury.
- [26] Einsle U. 1996. Copepoda: Cyclopoida Genera *Cyclops*, *Megacyclops*, *Acanthocyclops*. *Guides to the Identification of the Microinvertebrates of the Continental Waters of the*

- World, ed. H.J. Dumont. Amsterdam, The Netherlands: SPB Academic Publishing bv. 83.
- [27] Karaytuğ S. 1999. Copepoda:Cyclopoida Genera *Paracyclops*, *Ochridacyclops* and Key to the Eucyclopinae. Guides to the Identification of the Microinvertebrates of the Continental Waters of the World, 14, ed. H.J. Dumont. Backhuys Publishers, Leiden, 217pp.
- [28] Ueda H., Reid J. W. 2003. Copepoda:Cyclopoida Genera *Mesocyclops* and *Thermocyclops*. Guides to the Identification of the Microinvertebrates of the Continental Waters of the World, ed. H.J. Dumont. Backhuys Publishers, Leiden, 318pp.
- [29] Borutzky E.V. 1952. Fauna USSR. Crustacea, Volume III, No. 4. Freshwater Harpacticoida. Izdaniya Akademia Nauk SSSR, 50 [original in Russian but consulted for this work in the English translation by A. Mercado published in 1964 by the Israel Program for Scientific Translation, Jerusalem].
- [30] Damian-Georgescu A. 1970. Fauna Republicii Socialiste Romania, Crustacea, Copepoda, Harpacticoida, IV, 11. Academiei Republicii socialiste Romania. 248pp.
- [31] Wells J.B.J. 2007. An annotated checklist and keys to the species of Copepoda Harpacticoida (Crustacea). Zootaxa. 1568: 1-872.
- [32] Martin J.W., Davis G.E. 2001. An updated classification of the Recent Crustacea. Contr. Sci. Los Angeles No.39: i-x, 1-124.
- [33] Lang K. 1948. Monographie der Harpacticiden, 1-2: 1-1682. HÅkan Ohlsson, Lund.
- [34] Borutzky E.W. 1934. Zur Harpacticidenfuna des Kaukasus. Zool. Anzeig. 107: 263-269.
- [35] Kunz H. 1961. Beitrag zur Kenntnis der D'Arcythompsoniidae (Copepoda Harpacticoidea). Zool. Anzeig. 167(7-8): 275-280.
- [36] Chappuis P.A. 1953. Copépodes harpacticoides de Turquie. Bulletin de la Société d'histoire naturelle de Toulouse. 88: 355-361.
- [37] Dussart B.H., Defaye D. 1990. Répertoire mondial des Crustacés Copépodes des eaux intérieures. III. Harpacticoïdes. Crustaceana, Supplement 16: 1-364.