

Finnish Zoological and Botanical Publishing Board

Cirsium yildizianum (Asteraceae: Cynareae), a new species from East Anatolia, Turkey

Author(s): Turan Arabacı and Tuncay Dirmenci

Source: *Annales Botanici Fennici*, Vol. 48, No. 6 (2011), pp. 503-506

Published by: Finnish Zoological and Botanical Publishing Board

Stable URL: <https://www.jstor.org/stable/23728103>

Accessed: 02-12-2019 12:49 UTC

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at

<https://about.jstor.org/terms>



JSTOR

Finnish Zoological and Botanical Publishing Board is collaborating with JSTOR to digitize, preserve and extend access to *Annales Botanici Fennici*

Cirsium yildizianum (Asteraceae: Cynareae), a new species from East Anatolia, Turkey

Turan Arabaci^{1,*} & Tuncay Dirmenci²

¹⁾ İnönü University, Faculty of Science and Arts, Department of Biology, 44280 Malatya, Turkey
(*corresponding author's e-mail: tarabaci@inonu.edu.tr)

²⁾ Balıkesir University, Necatibey Education Faculty, Department of Biology Education, 10100 Balıkesir, Turkey

Received 17 May 2010, revised version received 9 Aug. 2010, accepted 10 Aug. 2010

Arabaci, T. & Dirmenci, T. 2011: *Cirsium yildizianum* (Asteraceae: Cynareae), a new species from East Anatolia, Turkey. — Ann. Bot. Fennici 48: 503–506.

Cirsium yildizianum Arabaci & Dirmenci sp. nova (Asteraceae: Cynareae) is described from East Anatolia, Turkey. It belongs to the section *Epitrachys* and is morphologically close to *C. macrobotrys* and *C. turkestanicum* s. lato. It differs from them mainly in its habit, distinctly pedunculate inflorescence, capitula number and the features of the phyllaries.

Cirsium is one of the largest genera within the family Asteraceae (Carduoideae: Cynareae/ Cardueae) and comprises approximately 250 species, distributed in Eurasia, North America, northern and eastern Africa, and growing mainly in wet wastelands and steppes (Boissier 1875, Davis & Parris 1975, Charadze 1976, Werner 1976, Petrak 1979, Kadereit & Jeffrey 2007). In Turkey, *Cirsium* is represented by 61 species (75 taxa), of which 20 (25 taxa) are endemic. Endemic and non-endemic species are mainly distributed in the E and NE Anatolia. The Turkish *Cirsium* species are classified in three sections: sect. *Epitrachys* (43 species), sect. *Cirsium* (17 species) and sect. *Cephalonoplos* (one species) (Davis & Parris 1975, Davis *et al.* 1988, Güner *et al.* 2000, Daşkın *et al.* 2006, Yıldız & Dirmenci 2008, Yıldız *et al.* 2009a, 2009b).

When the *Cirsium* species growing in Turkey were revised for the *Flora of Turkey*, some purple-flowered specimens (Sintenis 1890:3293,

Davis 24779 & 47412, Rechinger 15134) collected from the B7 and B8 squares were wrongly included in *C. macrobotrys* by Davis and Parris (1975). Our studies showed that these specimens are different from *C. macrobotrys*. In addition, during our revisionary studies of *Cirsium* in Turkey, in 2007–2008 we collected some material from the same squares and other parts of E Anatolia. These specimens were checked against the relevant literature (Boissier 1875, Davis & Parris 1975, Charadze 1976, Petrak 1979, Huber-Morath 1980, 1982, Sorger & Buchner 1983a, 1983b, Davis *et al.* 1988, Güner *et al.* 2000, Özhataç *et al.* 2009). Related specimens deposited in the herbaria ANK, BM, E, G, GAZI, HUB, ISTE, ISTF, ISTO, K, W and WU were also examined. It was concluded that the specimens collected from E Anatolia were representatives of a previously undescribed species in *Cirsium* sect. *Epitrachys* with affinities to *C. macrobotrys* and *C. turkestanicum* s. lato.




Fig. 1. *Cirsium yildizianum* (from the holotype). — A: Outer phyllary. — B: Median phyllary. — C: Inner phyllary. — D: Achene. — E: Median caudine leaf. — F: Basal leaf.

***Cirsium yildizianum* Arabacı & Dirmenci, sp. nova** (Figs. 1 and 2A–B)

Affinis *C. macrobotrys* Boiss. & *C. turkestanici* Petr. s. lato; ab *C. macrobotrys* capitulis 15–30 in medium usque ad apici caulis, (non 3–12 in apici caulis), inflorescentia racemosa vel breviter paniculata (non spica vel breviter racemosa), pedunculis (1–)3–8(–10) cm longis (non subsessilis), phyla exteriora 14–25 × 0.5–1.2 mm (non 20–45 × 1–2.5 mm), plerumque non excedens media (excedens media), phyllariis medianis oblongis ad lanceolatis (non ovatis) differt. Ab *C. turkestanici* s. lato biennis (non perennis), caulis 70–250 cm altus (non 40–100 cm), capitulis 15–30 in medium usque ad apici caulis, (non 3–7 in apici caulis), inflorescentia racemosa vel breviter paniculata (non breviter racemosa), pedunculis (1–)3–8(–10) cm longis (non sessilia vel brevissime pedunculata), phyllariis exterioris subulatis (non lanceolatis) differt.

TYPE: Turkey. B8 Muş: 4 km from Muş to Bingöl, 1380 m, wet places near field borders, 13.VIII.2007




Fig. 2. A–B: *Cirsium yildizianum* (type material). — A: Capitulum. — B: Inflorescence. — C: Inflorescence of *C. macrobotrys*.

Dirmenci 3567 & Arabacı (holotype INU; isotypes GAZI, HUB, ISTE).

ETYMOLOGY: The species is named after Prof. Dr. Bayram Yıldız, a well-known plant taxonomist in Turkey and also our Ph.D. supervisor.

Biennial. Stem 70–250 cm, capitulate from middle to apex, erect, stout, unwinged, longitudinally striate, pannose. Basal leaves 15–35 × 5–20 cm (except 3–10 cm petiole), oblong, pinnatisect, spinose-strigose above, setae 1–2.5 × 0.1–0.2 mm, adpressed, 5 or more per 2 mm square, otherwise glabrous; pannose beneath; lateral lobes 4–5 paired, 2–8 × 0.5–1 cm, lanceolate, acute to acuminate, apical spine 8–12 × 0.3–0.6 mm, moderately stout; stem leaves diminishing from base to inflorescence, median caudine leaves 7–12 × 4–6 cm, oblong to ovate, semiamplexicaul, pinnatisect to 1/4–1/5, 3–5 lobed, spinose-strigose above, setae 1–2.5 × 0.1–0.2 mm, 5 or more per 2 mm square, otherwise glabrous; pannose beneath;

lateral and terminal lobes ovate-triangular to lanceolate, acute, with apical spine $7–12 \times 0.2–6$ mm; upper caudine leaves to 10×6 cm, narrowly ovate, 2–3 lobed, lobes triangular, with $10–14 \times 0.5–0.8$ mm stout apical spine. Involucral leaves 8–15, 6–10 cm, linear-lanceolate, as long as or longer than involucre. Inflorescence raceme or shortly paniculate; capitula 15–30, erect, 1(–3) on each branch, $(20–)30–40 \times (20–)30–45$ mm, peduncle $(1–)3–8(–10)$ cm; involucres 20–30 × 30–35 mm, ovoid to globose; phyllaries 8–11 seriate, adpressed, glabrous below, arachnoid only at apex, yellowish; outer $14–25 \times 0.5–1.2$ mm, subulate, including erecto-patent to reflexed $2–8 \times 0.1–0.2$ mm apical spine, sometimes spinulose below apical spine; median $14–16 \times 1.4–1.6$ mm, oblong-lanceolate, abruptly narrowed in to prolongation, including recurved $1.5–2.5 \times 0.1–0.2$ mm apical spine; inner $18–24 \times 1–1.2$ mm, linear, including recurved $1–2 \times 0.1–0.2$ mm apical spine. Corolla purple, rarely ochroleucous, 18–25 mm, unequally 5 lobed to 1/6–1/4, shorter lobes 3–4 mm, longer lobes 5–6 mm; style 10–15 mm, exserted to 5 mm, shortly bilobed; filaments 5 mm, hairy, anthers 8–13 mm, acuminate. Achenes $6–6.5 \times 2–2.2$ mm, oblong, yellowish-brown, slightly asymmetric, compressed, umbo ca. 0.4 mm, ring yellow, narrow. Pappus long pluomose, 18–20 mm, stramineous. Flowering and fruiting from July to September.

Cirsium yildizianum appears to be endemic to E Anatolia and is an Irano-Turanian taxon. It grows in wet places near field borders, deep soiled areas, *Quercus* scrub openings and steppes at altitudes between 1380 and 2150 m.

Cirsium yildizianum is related to *C. macrobotrys* and *C. turkestanicum s. lato* in the foliar characteristics, but it can be easily distinguished from both of them by its stem, which capitulates from the middle to the apex, and by the racemose or shortly paniculate inflorescence, whereas *C. macrobotrys* and *C. turkestanicum* capitulate only at the apex of stem and the inflorescence is a spike or short-racemose (Table 1 and Fig. 2). In addition, *C. yildizianum* and *C. turkestanicum* are geographically isolated. The latter is distributed mainly in central Asia, from Tian-Shan to Pamir Mountains, Afghanistan, Turkmenistan, and NE Iran (Charadze 1976, Petrak 1979, Ghahremaninejad *et al.* 2005), whereas *C. yildizianum* is so far only recorded from E Anatolia in Turkey.

ADDITIONAL SPECIMENS EXAMINED: — *Cirsium yildizianum* (paratypes). Turkey. B7 Elazığ: Sivrice, between Gözeli and Kamişlık, 1500 m, deep soiled area, 2008 B. Yıldız 16964 & Arabacı; 38 km from Elazığ to Bingöl, 1700 m, *Quercus* scrub openings, 2007 Dirmenci 3508 & Arabacı; Erzincan: Sipikör, Sintenis (1890:3293) (G); B8 Erzincan: 85 km WSW Erzurum towards Tercan, 1956 Rechinger, 15134 (W); Palandöken Mount, 30–40 km from Çat to Erzurum, 2000–

Table 1. Morphological comparison of *Cirsium yildizianum*, *C. macrobotrys* and *C. turkestanicum s. lato*.

Species	<i>C. yildizianum</i>	<i>C. macrobotrys</i>	<i>C. turkestanicum s. lato</i>
Habit	biennial		
Stem	stem 70–250 cm, capitulate from middle to apex	biennial or shortly perennial, monocarpic stem 30–200(–300) cm, capitulate at apex	shortly perennial, monocarpic stem 40–100 cm, capitulate at apex
Inflorescence	raceme or shortly paniculate	spike or short-racemose	short-racemose
Capitula	15–30	(3–)5–12	3–4(–7)
Peduncle	(1–)3–8(–10) cm	sessile	sessile or very shortly pedunculate
Outer phyllary	generally not exceeding medians, subulate, apical spine 2–8 mm	exceeding medians, subulate, apical spine 5–10 mm	not exceeding medians, lanceolate, apical spine 2–4 mm
Median phyllary	14–16 mm, oblong-lanceolate, abruptly narrowed into prolongation	13–19 mm, ovate, abruptly narrowed into prolongation	oblong, suddenly attenuate
Inner phyllary	apical spine 1–2 × 0.1–0.2 mm	apical spine 2–3 × 0.1–0.2 mm	apical spine 1–2 × 0.1–0.2 mm
Pappus	18–20 mm	22–24 mm	21–25 mm

2150 m, in pasture, 1966 Davis 47412 (E, K); 32 km from Çat to Erzurum, 39°47'78''N, 41°05'00''E, 2150 m, 2007 Dirmenci 3529 & Arabaci; Bingöl: 15 km from Karlıova to Bingöl, 1950 m, steppe, 2007 Dirmenci 3516 & Arabaci; Muş: 24 km E of Muş, Davis 24779 (BM, E); C9 Hakkari: 15 km from Yüksekovalı to Hakkari, 1700–1800 m, 2007 Dirmenci 3571 & Fırat; 18 km from Şemdinli to Yüksekovalı, ca. 1700 m, Dirmenci 3580 & Fırat. — *C. macrobotrys*. **Turkey.** A7 Gümüşhane: Between Şiran and Alucra, Yıldız 16380 & Dirmenci; A8 Erzurum/Rize: Between İspir and Rize, Yıldız 16356 & Dirmenci; A9 Kars: Yağmurlu Mountain, Davis 32610 & Hedge (K, BM); B7 Sivas: 5 km west of İmraniye, Yıldız 16237 & Dirmenci; B8 Bayburt: Kop Mountain, Yıldız 16259 & Dirmenci; Erzurum: Kop Dağ pass, Furse 3835 (K); Palandöken Mountain, Yıldız 16272 & Dirmenci. — *C. turkestanicum* var. *turkestanicum*. **Afghanistan.** Sangcharak: Sufedsang, Koelz 13985 (W 1956: 594); Kabul: Paghamtales towards Kotal-i-Kotandar, A.Gilli 4158 (W 1963: 7419; type of *C. afghanicum*); in declibus orientalibus jugi Unai, Podlech 12593 (W 1967: 18475).

Acknowledgements

We would like to thank TÜBİTAK due to the financial support our researches (Project no. 106T167), and SYNTHESYS Project (AT-TAF58 & GB-TAF3087) which is financed by European Community Research Infrastructure Action under the FP6 “Structuring the European Research Area” Programme for financial support our studies in BM, E, K and W and the curators of these herbaria, whose gave us permission to examine the specimens. Also, our thanks to curators of the herbaria ANK, G, GAZI, HUB, ISTE, ISTF, ISTO and WU.

References

- Boissier, E. 1875: *Flora Orientalis*, 3: 523–553. — Genevae et Basileae.
- Charadze, A. L. 1976: *Cirsium* Mill. — In: Bobrov, E. G. & Cherepanov, S. K. (eds.), *Flora of the U.S.S.R.*, 28: 63–270. Israel Program for Scientific Translations, Jerusalem.
- Daşkin, R., Yılmaz, Ö. & Kaynak, G. 2006: Presence of *Cirsium eriophorum* (L.) Scop. (Asteraceae) in Turkey. — *Turkish J. Bot.* 30: 1–5.
- Davis, P. H. & Parris, S. B. 1975: *Cirsium* Mill. — In: Davis, P. H. (ed.), *Flora of Turkey and the East Aegean Islands*, vol. 5: 370–412. Edinburgh Univ. Press, Edinburgh.
- Davis, P. H., Tan, K. & Mill, R. R. (eds.) 1988: *Flora of Turkey and the East Aegean Islands*, Suppl. 1, 10: 164–165. — Edinburgh Univ. Press, Edinburgh.
- Ghahremannejad, F., Joharchi, M. & Vitek, E. 2005: New plant records for Khorasan province, Iran. — *Ann. Naturhist. Mus. Wien* 106: 255–293.
- Güner, A., Özhatay, N., Ekim, T. & Başer, K. H. C. (eds.) 2000: *Flora of Turkey and the East Aegean Islands*, Suppl. 2, 11: 161–163. — Edinburgh Univ. Press, Edinburgh.
- Huber-Morath, A. 1980: Ergänzungen zu PH. Davis' Flora of Turkey and the East Aegean Islands 1–6. — *Candollea* 35: 569–608.
- Huber-Morath, A. 1982: Ergänzungen zu PH. Davis' Flora of Turkey and the East Aegean Islands. — *Candollea* 39: 323–344.
- Kadereit, J. W. & Jeffrey, C. (eds.) 2007: Flowering plants. Eudicots: Asterales. — In: Kubitzki, K. (ed.), *The families and genera of vascular plants*, vol. 8: 132. Springer-Verlag, Berlin–Heidelberg.
- Özhatay, N., Kültür, S. & Aslan, S. 2009: Check-list of additional taxa to the supplement flora of Turkey IV. — *Turkish J. Bot.* 33: 191–226.
- Petrak, F. 1979: *Cirsium* Mill. — In: Rechinger, K. H. (ed.), *Flora Iranica* vol 139a: *Compositae III – Cynareae*: 231–280. Akademische Druck- und Verlagsanstalt, Graz, Austria.
- Sorger, F. & Buchner, P. 1983a: Beiträge zur Flora der Türkei III. — *Phyton* 23: 221–245.
- Sorger, F. & Buchner, P. 1983b: Beiträge zur Flora der Türkei III. — *Linzer Biol. Beiträge* 14: 157–208.
- Werner, K. 1976: *Cirsium* Mill. — In: Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M. & Webb, D. A. (eds.), *Flora Europaea*, vol. 4: 232–242. Cambridge Univ. Press, Cambridge.
- Yıldız, B. & Dirmenci, T. 2008: A new species of *Cirsium* section *Epitrachys* (Asteraceae: Cardueae) from Turkey. — *Bot. J. Linn. Soc.* 158: 669–673.
- Yıldız, B., Dirmenci, T. & Arabaci, T. 2009a: *Cirsium handianae* (Asteraceae), a new species from Turkey. — *Annales Botanici Fennici* 46: 239–243.
- Yıldız, B., Dirmenci, T. & Arabaci, T. 2009b: A new record for the flora of Turkey: *Cirsium candelabrum* Griseb. (*Cirsium* Sect. *Cirsium*, Asteraceae, Cynareae). — *Turkish J. Bot.* 33: 47–51.