

# Seasonal Variations of Phytoplankton Community in Relation to Some Physical and Chemical Parameters in a Temperate Eutrophic Reservoir, Turkey

Kemal Çelik<sup>1\*</sup>, Tuğba Ongun Sevindik<sup>2</sup>

<sup>1</sup>Balıkesir University, Faculty of Arts and Science, Department of Biology, 10145, Balıkesir, Turkey

<sup>2</sup>Sakarya University, Faculty of Arts and Science, Department of Biology, 54187, Adapazarı, Turkey

Received Revised Accepted

## Abstract

The Çaygören Reservoir is fed by the Simav Stream and the maximum inflow ( $1300 \text{ m}^3 \text{ sec}^{-1}$ ) occurred in spring and the minimum (about  $5.2 \text{ m}^3 \text{ sec}^{-1}$ ) occurred in fall. It has an annual mean water capacity of  $392 \text{ hm}^3$  and a total volume of  $142.57 \text{ hm}^3$ . A total of 192 taxa in 9 divisions were identified. *Cyclotella meneghiniana* Kützing, *Stephanodiscus neoastraea* Hakansson and Hickel of Bacillariophyta, *Gloeotila subconstricta* (G.S.West) Printz of Chlorophyta, *Mugeotia* sp. of Streptophyta, *Cryptomonas pyrenoidifera* Geitler, *Plagioselmis nannoplantica* (H.Skuja) G.Novarino, I.A.N.Lucas and S.Morrall of Cryptophyta, *Aphanocapsa holsatica* (Lemmermann) G.Cronberg and J.Komárek, *Aphanothece clathrata* West and G.S.West and *Planktothrix* sp. of Cyanobacteria dominated phytoplankton at least for one season during the observation period. Species of Cryptophyta dominated phytoplankton during the winter, while Chlorophyta and Streptophyta species were dominant in the fall. Bacillariophyta species dominated phytoplankton in the spring and Cyanobacteria were dominant in the summer. The maximum phytoplankton biomass and abundance ( $106.5 \text{ mg L}^{-1}$ ;  $273154 \text{ individual M}^{-3}$ ) were recorded in summer 2008 at the third station and the minimum biomass and abundance ( $0.23 \text{ mg L}^{-1}$ ;  $799 \text{ individual M}^{-3}$ ) were recorded in winter 2007 at the second station. The canonical correspondence analysis (CCA) and correlation results showed that water temperature, transparency, phosphate, oxidation-reduction potential and water discharge had significant effects (Monte Carlo test,  $p < 0.05$ ) on the dynamics of dominant phytoplankton of the eutrophic Çaygören Reservoir.

**Keywords:** Phytoplankton, Temperate Eutrophic Reservoir, Water Discharge, Water Quality Parameters, CCA.

## 1. Introduction

Çaygören reservoir was built between 1965 and 1968 for the purpose of irrigation and hydropower generation. It is an important source of irrigation water for the towns of Sındırgı and Bigadiç in the province of Balıkesir, Turkey. It is used for flood control as its source stream (Simav Stream) sometimes reaches about  $1500 \text{ m}^3 \text{ sec}^{-1}$  debt. The Çaygören Reservoir has a total length of 4.6 km, a surface area of  $8.15 \text{ km}^2$  and a maximum depth of 53.5 m. The purpose of the present study is to determine the environmental variables responsible for the seasonal variations of the abundance, species composition and the biomass of the phytoplankton community of the temperate eutrophic Çaygören Reservoir.

Phytoplankton communities play an important role in aquatic ecosystems as they produce food and oxygen, which supports all other life forms (Fathiet al., 2001). Knowledge on their abundance and community composition provided further understanding of ecological interactions in aquatic ecosystems. The seasonal dynamics of phytoplankton have been investigated worldwide (Taveriniet al., 2009; Jeppesen et al., 2011; Elliott, 2012; Caroni et al., 2012; Feuchtmayret et al., 2012).

In northern temperate lakes, phytoplankton succession is largely determined by the seasonal cycles of physical, chemical and biological factors, the relative importance of which varies with the different periods of the year (Tiina et al., 2011). The spatial distribution of phytoplankton in lakes is highly heterogeneous. This heterogeneous distribution is mostly attributed to wind events,

\* Corresponding author. e-mail: kcelik@balikesir.edu.tr.

mixing and the contrasting gradients in the light and nutrient concentrations (Reynolds et al., 2002).

Considering the complexity of phytoplankton dynamics, studying the spatial and seasonal distribution of species and their relationships with the physicochemical parameters can give insights into understanding factors responsible for their dynamics. Arhonditsis et al. (2004) stated that to determine which factors are effective on spatial and temporal distribution of phytoplankton, the system under study should have been sampled for at least two years.

The underwater light climate seems to be one of the selective environmental factors that strongly influence the species composition and biomass of phytoplankton in lakes. Water transparency has received much attention lately because of the self-limitation of photosynthesis imposed by phytoplankton. In turbid environments, algal species with gas vesicles can either move down to avoid the high light intensity at the water surface, or float up when underwater light conditions are poor (Pérez et al., 2007).

Nutrient limitation imposes compositional changes on the phytoplankton community. According to resource competition theory, nitrogen-fixing algae should dominate lakes when nitrogen is limiting. Assuming that phosphorus and nitrogen limit many algae in lakes, blue-green algae should dominate lakes when N:P ratios are low but other types should dominate when they are high (Chaffin et al., 2011).

For a better understanding of the processes affecting phytoplankton dynamics, it is important to study the linkage between changes in environmental variables and phytoplankton abundance, biomass and community composition (George et al., 2004). Multivariate statistical techniques have been proved to be useful for understanding interactions between the ecological factors and plankton communities in aquatic ecosystems (Kruk et al., 2002).


Although few studies have been published on the Çaygören Reservoir (Sevindik, 2010; Sevindiket et al., 2011), the present study is the first attempt to describe the seasonal and spatial distribution of environmental variables and their relations with the phytoplankton composition in the temperate eutrophic Çaygören Reservoir using Canonical Correspondence Analysis (CCA) and Pearson's correlation analysis.

## 2. Materials and Method

### 2.1. Study Area

The Çaygören Reservoir is located at 39° 17' 24" N; 28° 19' 16" E, 55 km southeast of Balıkesir, Turkey (Fig. 1). It lies at 273 m above the sea level

and has a maximum depth of 28 m, a length of 4.6 km and a surface area of 9 km<sup>2</sup>. The reservoir is fed by the Simav Stream. Its construction started in 1971 and it is used for irrigation and power generation (State Water Works, 2005).



**Figure 1.** The map of the Çaygören Reservoir and the location of sampling stations (I think this should be placed under the figure above this paragraph or the figure should be moved to above this caption)

### 2.2. Sampling Procedure and Chemical Analysis

Water sampling was carried out monthly from February 2007 to January 2009 for measurements of physical, chemical and biological parameters. Samples were collected vertically at 5 m intervals using a Kemmerer water sampler. Specific Conductivity (SC), pH, Oxidation-Reduction Potential (ORP) and water Temperature (T) were measured at 1 m intervals using a YSI multi probe. Water transparency was measured using a Secchi disk.

Concentrations of phosphate (PO<sub>4</sub>), nitrate-nitrogen (NO<sub>3</sub>-N) and ammonium-nitrogen (NH<sub>4</sub>-N) were determined spectrophotometrically in samples collected from 1, 5 and 15 meters according to the standard methods (APHA, 1995). Water Discharge (WD) data were obtained from the State Water Works.

For the purpose of minimizing the errors, calibration of apparatus, running of blank and sample at known concentration, measurements in replicate were performed in the laboratory. The accuracy and precision of the used analytical methods were checked by means of standard samples, which were assayed with each series of samples (Table 1). A multipoint calibration of the YSI multi probe was done one day prior to the sampling. Zero and span checks were made regularly as the basic quality assurance procedure for analysis.

**Table 1.** Mean  $\pm$  Standard deviation (SD) physical and chemical water characteristics for water quality parameters in the Çaygören Reservoir

Parameter	Depth	Control	St. 1	St. 2	St.3
Tmp (°C)	1 m	15.2 $\pm$ 6.35	14 $\pm$ 6.8	14 $\pm$ 6.3	15 $\pm$ 6.3
	10 m	16.1 $\pm$ 7.71	15 $\pm$ 7.8	14 $\pm$ 6.4	15 $\pm$ 6.45
	15 m	13.9 $\pm$ 7.65	14.6 $\pm$ 7.1	13.9 $\pm$ 6.39	13.9 $\pm$ 6.38
Cond (mS cm <sup>-1</sup> )	1m	0.44 $\pm$ 0.089	0.43 $\pm$ 0.09	0.44 $\pm$ 0.07	0.43 $\pm$ 0.08
	10 m	0.46 $\pm$ 0.081	0.44 $\pm$ 0.08	0.24 $\pm$ 0.04	0.24 $\pm$ 0.07
	15 m	0.45 $\pm$ 0.0074	0.44 $\pm$ 0.075	0.23 $\pm$ 0.05	0.23 $\pm$ 0.06
pH	1 m	9.9 $\pm$ 0.59	9.8 $\pm$ 0.6	9.78 $\pm$ 0.61	9.73 $\pm$ 0.63
	10 m	9.9 $\pm$ 0.579	9.7 $\pm$ 0.59	9.5 $\pm$ 62	9.87 $\pm$ 61
	15 m	9.8 $\pm$ 0.61	9.75 $\pm$ 0.58	10.9 $\pm$ 5.91	10.8 $\pm$ 5.71
ORP (mV)	1 m	102 $\pm$ 43	102 $\pm$ 42	101 $\pm$ 48	101 $\pm$ 43
	10 m	103 $\pm$ 44.5	102 $\pm$ 45	104 $\pm$ 31	104 $\pm$ 32
	15 m	100 $\pm$ 43.12	101 $\pm$ 43	103 $\pm$ 34	103 $\pm$ 33
PO <sub>4</sub> (mg L <sup>-1</sup> )	1 m	0.023 $\pm$ 0.009	0.022 $\pm$ 0.008	0.02 $\pm$ 0.002	0.02 $\pm$ 0.0012
	10 m	0.024 $\pm$ 0.009	0.024 $\pm$ 0.01	0.034 $\pm$ 0.02	0.031 $\pm$ 0.013
	15 m	0.024 $\pm$ 0.01	0.023 $\pm$ 0.011	0.031 $\pm$ 0.0079	0.03 $\pm$ 0.0071
NO <sub>3</sub> (mg L <sup>-1</sup> )	1 m	0.19 $\pm$ 0.051	0.2 $\pm$ 0.05	0.21 $\pm$ 0.05	0.23 $\pm$ 0.045
	10 m	0.21 $\pm$ 0.043	0.2 $\pm$ 0.04	0.18 $\pm$ 0.04	0.19 $\pm$ 0.038
	15 m	0.19 $\pm$ 0.041	0.18 $\pm$ 0.039	0.21 $\pm$ 0.045	0.23 $\pm$ 0.042
NH <sub>4</sub> (mg L <sup>-1</sup> )	1m	0.048 $\pm$ 0.0023	0.05 $\pm$ 0.002	0.025 $\pm$ 0.0012	0.042 $\pm$ 0.0011
	10 m	0.071 $\pm$ 0.0051	0.07 $\pm$ 0.005	0.023 $\pm$ 0.0011	0.041 $\pm$ 0.0013
	15 m	0.072 $\pm$ 0.0025	0.068 $\pm$ 0.002	0.043 $\pm$ 0.0021	0.043 $\pm$ 0.0012
TSS (mg L <sup>-1</sup> )	1 m	15.5 $\pm$ 6.56	16 $\pm$ 6.3	14 $\pm$ 4.5	14 $\pm$ 4.5
	10 m	17 $\pm$ 7.69	17 $\pm$ 7.7	14.6 $\pm$ 5.7	15.6 $\pm$ 4.7
	15 m	16.8 $\pm$ 7.25	16 $\pm$ 7.2	15.9 $\pm$ 6.5	15.9 $\pm$ 4.5

### 2.3. Phytoplankton Sampling and Analysis

In the field, samples for phytoplankton were collected from 1, 5 and 15 meters and placed in 250 ml bottles and fixed with Lugol's solution. In the laboratory, the samples were first agitated, then poured into 50 ml graduated cylinders and were allowed to settle for 24 hours. At the end of the settling period, 45 ml of water was aspirated from each graduated cylinder and the remaining 5 ml was poured into a small glass vial for microscopic analysis. Enumeration and identification of phytoplankton were performed using a Palmer-Maloney counting cell and an Olympus BX 51 compound microscope equipped with water immersion lenses (40X and 60X magnifications) and a phase-contrast attachment.

Phytoplankton species were identified according to Huber-Pestalozzi (1941; 1950; 1969; 1972; 1982; and 1983), Bourrelly (1968), Krammer and Lange-Bertalot (1986; 1991; and 1999), Komarek and Anagnostidis (1986; 1989; 1999; and 2008), Anagnostidis and Komarek (1988), Round et al. (1990), Sims (1996) and John et al. (2003). Phytoplankton biomass was calculated from the biovolume data, assuming a specific gravity of one (Edmondson, 1971). Biovolume was calculated from cell numbers and cell size measurements (Sun and Liu, 2003).

### 2.4. Statistical Analysis

The CCA was used to determine the relationships between the dominant phytoplankton taxa and the environmental variables. The significance of environmental variables on the dominant taxa was determined with Monte Carlo tests. The CCA and Monte Carlo tests were performed using the CANOCO v.4.5 program (ter Braak and Smilauer, 2002). The relationships between the physicochemical variables and the dominant phytoplankton taxa were further analyzed using the Pearson's correlation coefficients.


The statistical differences in the total phytoplankton abundance and biomass were determined using an ANOVA test. The ANOVA and the Pearson's correlation coefficients were calculated using SAS statistical software. Data were log transformed before the statistical analysis to obtain normal distribution (SAS Institute, 2003).

## 3. Results


### 3.1. Physico-Chemical Parameters

The maximum (1300 m<sup>3</sup> s<sup>-1</sup>) and minimum (5.2 m<sup>3</sup> s<sup>-1</sup>) inflows (WD) were recorded in April 2007 and September 2007, respectively (Fig. 2). Secchi disk depth ranged from 0.3 m to 1.5 m at St.1 and it


ranged from 0.6 m to 1.9 m at St.2 and St.3 (Fig. 3). Water temperature ranged from 4.5°C to 27.6 °C at all stations. Maximum surface water temperature values were measured in June and July and minimum values were measured in February (Fig. 4).



**Figure 2.** The seasonal variations in the water discharge ( $m^3 s^{-1}$ ) of the Çaygören Reservoir






**Figure 3.** The seasonal variations in the Secchi disk depth (m) of the Çaygören Reservoir




**Figure 4.** The seasonal variations in the water temperature (°C) of the Çaygören Reservoir.

Specific conductivity ranged from 0.3  $mS cm^{-1}$  to 0.6  $mS cm^{-1}$  at all stations and it was lower in the winter than the other seasons (Fig. 5). pH ranged from 7.4 to 11.6 from February 2007 to September 2008 at all stations. From November 2008 to January 2009, pH fluctuated between 7.7 and 11 (Fig. 6).




**Figure 5.** The seasonal variations in specific conductivity ( $mS cm^{-1}$ ) of the Çaygören Reservoir




**Figure 6.** The seasonal variations in the pH of the Çaygören Reservoir

ORP ranged from 2 mV to 219.5 mV at all stations and it was lower in the summer than the other seasons (Fig. 7). PO<sub>4</sub> concentrations ranged from 0.005 mg L<sup>-1</sup> to 0.06 mg L<sup>-1</sup>, oscillating around 0.02 mg L<sup>-1</sup> throughout the study period, except for a peak of 0.04 mg L<sup>-1</sup> in October 2008 at the first station and another one of 0.06 mg L<sup>-1</sup> in November 2007 at the second station (Fig. 8).




**Figure 7.** The seasonal variations in the oxidation-reduction potential (mV) of the Çaygören Reservoir




**Figure 8.** The seasonal variations in the phosphate (mg L<sup>-1</sup>) of the Çaygören Reservoir

NO<sub>3</sub>-N concentrations ranged from 0.055 mg L<sup>-1</sup> to 0.3 mg L<sup>-1</sup> at all stations. A decline of about 0.05 mg L<sup>-1</sup> in NO<sub>3</sub>-N occurred in the spring and fall at all stations during the study period (Fig.9). NH<sub>4</sub>-N concentrations ranged from 0.005 mg L<sup>-1</sup> to 0.017 mg L<sup>-1</sup> at the first and second stations and they ranged from 0.001 mg L<sup>-1</sup> to 0.02 mg L<sup>-1</sup> at the third station (Fig. 10).




**Figure 9.** The seasonal variations in nitrate-nitrogen (mg L<sup>-1</sup>)



**Figure 10.** The seasonal variations in the ammonium-nitrogen (mg L<sup>-1</sup>) of the Çaygören Reservoir


**3.2. Phytoplankton Species and Biomass**

A total of 192 taxa in nine major taxonomic categories were identified. During the winter, *Plagioselmis nannoplantica* (H.Skuja) G.Novarino, I.A.N.Lucas and S.Morrall, Cryptophyta; 10% of the total biomass) dominated phytoplankton. In the spring, *Cyclotella meneghiniana* Kützing, (Bacillariophyta; 35% of the total biomass) and *Stephanodiscus neoastraea* Hakansson and Hickel (Bacillariophyta; 32% of the total biomass) were dominant. During the summer, *Planktothrix* sp. (Cyanobacteria; 33% of the total biomass), *Aphanocapsa holsatica* (Lemmermann) Cronberg (Cyanobacteria; 12.5% of the total biomass) and *Aphanothece clathrata* West and G.S.West (Cyanobacteria; 30% of the biomass) dominated phytoplankton. In the fall, *Gloetila subconstricta* (G.S. West) Printz (Chlorophyta; 10% of the total biomass) and *Mougeotia* sp. (Streptophyta; 14% of the total biomass) were dominant in the Çaygören Reservoir (Fig. 11).




**Figure 11.** The percent biomass distribution of the dominant phytoplankton taxa in the Çaygören Reservoir

The maximum phytoplankton biomass was recorded in winter 2007 (78 mg L<sup>-1</sup> at the first station, 99 mg L<sup>-1</sup> at the second station and 106.5 mg L<sup>-1</sup> at the third station) and the lowest biomass was recorded in December 2007 (0.31 mg L<sup>-1</sup> at the first and the third stations and 0.23 mg L<sup>-1</sup> at the second station; Fig. 12). The phytoplankton biomass was significantly different among the seasons (F=104, P<0.001), but not among the sampling stations (F=0.65, P>0.01).



**Figure 12.** The seasonal distribution of the total phytoplankton biomass (g L<sup>-1</sup>) in the Çaygören Reservoir

The maximum phytoplankton abundance was measured in summer 2008 because of optimum water temperature and sufficient light and the minimum abundance was recorded in winter 2007 because of low water temperature and insufficient light. The seasonal variations in phytoplankton abundance during 2007-2009 are presented in Fig. 13. The differences in the phytoplankton abundance were significant among the seasons ( $F=64$ ,  $P<0.001$ ), but not among the sampling stations ( $F=0.39$ ,  $P>0.05$ ).



**Figure 13.** The seasonal dynamics of the total phytoplankton abundance (cell L<sup>-1</sup>) in the Çaygören Reservoir

**3.3. Statistical Analysis of Phytoplankton Species, Biomass and Physico-Chemical Parameters**


In the Çaygören Reservoir, from CCA analysis, the first and second axes of CCA explained 77.2% of the total variance in the dominant phytoplankton taxa-environment relationships (eigenvalues, 0.8 and 0.55). The third and fourth axes explained 22.3% of the total variance (eigenvalues, 0.378 and 0.016). Table 2 shows the results of the Monte Carlo tests for the significance of the physicochemical parameters in order of the variance they explain. According to these results, water temperature, water discharge, Secchi disk transparency, oxidation-reduction potential and phosphate had significant effects on the dynamics of the phytoplankton ( $p<0.05$ ).

**Table 2.** The variance explained by each variable in the Çaygören Reservoir

Variable	Variable number	Variance explained	P	F
T (°C)	1	0.72	0.002*	7.91
WD (m <sup>3</sup> s <sup>-1</sup> )	9	0.53	0.002*	7.57
Secc. (m)	2	0.29	0.032*	3.02
Scand. (mS cm <sup>-1</sup> )	3	0.07	0.271	1.110
NO <sub>3</sub> -N (mg L <sup>-1</sup> )	6	0.02	0.382	0.154
ORP (mV)	4	0.26	0.038*	2.86
pH	5	0.05	0.584	0.261
NH <sub>4</sub> -N (mg L <sup>-1</sup> )	7	0.03	0.714	0.571
PO <sub>4</sub> (mg L <sup>-1</sup> )	8	0.25	0.039*	2.85

\*significant at 0.05 level.

The first axis of CCA was positively related to T, SC, NH<sub>4</sub>-N and PO<sub>4</sub> and it was negatively related with Secchi disk transparency, NO<sub>3</sub>-N, ORP, WD and pH. The second axis was positively related to Secchi disk transparency, NO<sub>3</sub>-N, ORP, pH and NH<sub>4</sub>-N and it was negatively related with T, SC, PO<sub>4</sub> and WD (Fig. 14).



**Figure 14.** The diagram of Canonical Correspondence Analysis (CCA) showing the relationships between the environmental variables and the dominant phytoplankton taxa in the Çaygören Reservoir. Abbreviation for species: Cycmen, *Cyclotella meneghiniana*; Stepneo, *Stephanodiscus neoastraea*; Glosub, *Gloeotilla subconstricta*; Aphhol, *Aphanocapsa holsatica*; Plankt, *Planktothrix* sp.; Cryptpy, *Cryptomonas pyrenoidifera*; Aphanotc, *Aphanothece clathrata*; Mougeotia sp.; Pnan, *Plagioselmis nannoplantctica*.

Fig. 14 shows the relationships between environmental variables and the dominant phytoplankton taxa. The distribution of cyanobacteria, *Planktothrix* sp., *A. holsatica* and *A. clathrata*, along the positive side of the first axis of CCA diagram, reflected their occurrence at high temperature. *G. subconstricta* (Chlorophyta) and *Mougeotia* sp. (Streptophyta) were located on the

positive side of the first axis and they were negatively related with water transparency and  $\text{NO}_3\text{-N}$ . The cryptophytes, *C. pyrenoidifera* and *P. nanoplantica* were located on the positive side of the second axis and they were negatively related with T. The diatoms, *C. meneghiniana* and *S. neoastreae* were located on the negative side of the second axis and they were positively related to WD.

**Table 3.** The Pearson's correlation coefficients between the physicochemical parameters and the dominant phytoplankton taxa in the Çaygören Reservoir between 2007 and 2009

	T	Secc	SC	ORP	PH	$\text{NO}_3$	$\text{NH}_4$	$\text{PO}_4$	WD
T	1	-0.5	0.6*	-0.6*	0.4	-0.6*	0.6*	-0.6*	0.1
Secc	-0.5	1	0.01	0.2	0.1	0.01	0.01	0.2	0.4
SC	0.6*	0	1	-0.8*	-0.5*	0.1	0.2	0.8*	0.4
ORP	-0.6*	0.2	-0.8*	1	0.5*	-0.8*	-0.4	-0.91*	0
pH	0.4	0.1	-0.5*	0.5	1	-0.5*	-0.1	0.5*	-0.1
$\text{NO}_3$	-0.6*	0	0.1	-0.8	-0.5*	1	0.2	-0.8*	0.4
$\text{NH}_4$	0.6*	0	0.2	-0.4	-0.1	0.2	1	-0.4	-0.1
$\text{PO}_4$	-0.6*	0.2	0.8*	-0.91*	0.5*	-0.8*	-0.4	1	0
WD	0.13	0.4	0.4	0.01	-0.1	0.4	-0.1	0.1	1
<i>Cycmen</i>	-0.2	0.2	0.1	0.2	0.2	0.1	-0.3	0.2	0.7*
<i>Stepneo</i>	-0.1	0.3	0.1	0.2	0.2	0.1	-0.3	0.2	0.7*
<i>Glosub</i>	-0.1	-0.6*	0.2	0.1	0.6*	0.2	-0.2	0.9*	-0.3
<i>Mougeot</i>	0.3	-0.5*	0.2	0.1	0.4	0.2	0	0.9*	-0.4
<i>Aphhol</i>	0.5*	-0.1	-0.2	-0.3	-0.4	-0.2	0.2	-0.3	0
<i>Aphanotc</i>	0.5*	0.1	0	-0.3	-0.6*	0	0.1	-0.3	0.2
<i>Plankt</i>	0.5*	0.2	-0.4	-0.4	-0.6*	-0.4	0.5*	-0.4	-0.3
<i>Cryptpy</i>	-0.7*	0.4	-0.5*	0.3	0.4	-0.45	-0.2	-0.42	-0.3
<i>Pnan</i>	-0.7*	0.4	-0.6*	0.3	0.41	-0.36	-0.2	-0.41	-0.4

\*significant at 0.05 level.

Abbreviation: *Cycmen*, *Cyclotella meneghiniana*; *Stepneo*, *Stephanodiscus neoastreae*; *Glosub*, *Gloeotila subconstricta*; *Aphhol*, *Aphanocapsa holsatica*; *Plankt*, *Planktothrix* sp.; *Pnan*, *Plagioselmis nanoplantica*; *Cryptpy*, *Cryptomonas pyrenoidifera*; *Aphanotc*, *Aphanothece clathrata*; *Mougeot*, *Mougeotia* sp.

There were significant correlations between most of the water quality parameters (Table 3). Significant positive correlations were observed between T and SC, T and  $\text{NH}_4$ ; pH and ORP, pH and  $\text{PO}_4$ ; ORP and  $\text{PO}_4$ , while negative correlations were obtained between T and  $\text{PO}_4$ , T and  $\text{NO}_3\text{-N}$ , T and ORP; SC and  $\text{PO}_4$ , SC and pH, SC and ORP; ORP and pH, ORP and  $\text{NO}_3\text{-N}$ ;  $\text{NO}_3\text{-N}$  and  $\text{PO}_4$ .

#### 4. Discussion

##### 4.1. Statistical Analysis and Interpretation of Physico-Chemical Parameters

The results of the present study showed that water temperature was negatively correlated with  $\text{NO}_3\text{-N}$  and it was positively correlated to  $\text{NH}_4\text{-N}$ . The significant positive correlation between ammonium concentrations and water temperature could probably be attributed to the intensified ammonification of  $\text{NO}_3\text{-N}$  with the increased water temperature (Liikanen and Martikainen, 2003). Ammonium is often used preferentially by phytoplankton over nitrate when both substrates are available in the water column (Stolte and Riegman, 1996). The negative correlation between the nitrate

There were significant correlations between the following the physicochemical variables and the dominant taxa:

Secchi disk transparency and *G. subconstricta*, Secchi disk transparency and *Mougeotiasp.*; pH and *G. subconstricta*, pH and *A. clathrata*;  $\text{NH}_4\text{-N}$  and *Planktothrix* sp.; WD and *C. meneghiniana* and WD and *S. neoastreae* (Table 3).

concentrations and water temperature may also indicate the effective consumption of the winter stock of nitrate by phytoplankton during the summer blooms in the Çaygören Reservoir (Temponeraset al., 2000).

ORP and  $\text{PO}_4$  were negatively correlated. ORP is the most important factor influencing the exchange process of phosphorus between the water and sediments (Li et al., 2010). It is well known that a decrease of ORP results in the deoxidization of metal-oxides, which might lead to a release of  $\text{PO}_4$ , whereas a rise of ORP helps to cause more  $\text{PO}_4$  to be absorbed (House and Deniso, 2000). It could be concluded that high  $\text{PO}_4$  concentrations in the summer were attributed to the transfer of  $\text{PO}_4$  from the bottom layers due to low ORP values in the Çaygören Reservoir.

Raised temperatures stimulate the overall mineralization and thereby liberate organic-bound phosphorus into the sediment pore water. In addition to this direct effect, increased microbial activity lowers the redox potential in the surface sediment, which may induce release of Fe-bound phosphorus (Wilhelm and Adrian, 2008).



There is a positive correlation between specific conductivity and  $PO_4$ . Specific conductivity is often considered as parameter showing the degree of nutrient loading (Parinetet *et al.*, 2004). Intensive agriculture has been practiced in the drainage basin of the Çaygören Reservoir. Agricultural nonpoint sources are a major contributing factor to surface water eutrophication worldwide.

Secchi disk transparency and water temperature were negatively correlated. In standing water bodies, turbidity increases with nutrient levels, which stimulate phytoplankton growth, especially during warm seasons (Scheffer and van Nes, 2007). Low Secchi disk depths in the summer timing are probably due to high abundance of phytoplankton. The eutrophic Çaygören Reservoir is dominated by Cyanobacteria in summer. Cyanobacteria strongly absorb light causing reduced water transparency (LaBounty, 2008).

Nitrate was negatively correlated with pH. It has long been known that the dense populations of phytoplankton deplete the carbon dioxide present in natural waters, resulting in the rise of pH. Jones-Lee and Lee (2005) state that algae take up nitrate and  $CO_2$  from water, causing the increase of pH during daylight in eutrophic lakes. The higher pH values found in the Çaygören Reservoir must accordingly have been a result of the depletion of free  $CO_2$  in the water due to high rate of photosynthesis.

pH is the master variable in the chemistry of aquatic systems and it affects the kinetics of nutrient uptake and controls the chemical species of most of the nutrient ions that algae require. Carbon fixation as a consequence of photosynthetic activity can displace the carbon dioxide-bicarbonate-carbonate equilibrium that is the most common pH-buffering mechanism in freshwater systems. Photosynthesis thus tends to increase pH in lakes (Haande *et al.*, 2011).

#### 4.2. Statistical Analysis and Interpretation of Phytoplankton and Physico-Chemical Parameter Relationships

In the CCA diagram, the cyanobacteria, *Planktothrix* sp., *A. clathrata* and *A. holsatica* occurred near  $NH_4$ -N and water temperature vectors. *Planktothrix* sp. was the most abundant cyanobacterium that dominated phytoplankton in the summer in the Çaygören Reservoir. The timing of *Planktothrix* bloom in the Çaygören Reservoir appears to be related to high temperature of the eutrophic condition of the reservoir. Temperature is one of the most important factors affecting the biology of phytoplankton species by controlling the rate of enzymatic reactions within the cells. In addition, temperature also regulates the multiplication rate and standing stock of natural phytoplankton populations (Niu *et al.*, 2011).

Padisak *et al.* (2009) state that small colonial non-N-fixing cyanobacteria prefer well-mixed environments at high water temperatures. The high nutrient levels in the Çaygören Reservoir probably

accounted for the development of this cyanobacterium during the summer time.

Water temperature had significant correlations with *A. clathrata* and *A. holsatica*. Komarek and Anagnostidis (1999) point out that *A. clathrata* and *A. holsatica* prefer eutrophic waters. *A. clathrata* and *A. holsatica* are widely collected in eutrophic Turkish lakes during the summer (Sevindiket *et al.*, 2010).

*C. meneghiniana* (Bacillariophyta) and *S. neoastrea* (Bacillariophyta) dominated phytoplankton in the spring. Diatoms are usually common during cooler or windier conditions in freshwater lakes (Munawar and Munawar, 1986). These species were occasionally abundant at the shallower first station. Although *Cyclotella* and *Stephanodiscus* species are widely collected in freshwater phytoplankton, some of them are also benthic (Hutchinson, 1967). They might have been drifted from the bottom due to wind-driven water turbulence at this shallow station.

In the CCA diagram, *S. neoastrea* and *C. meneghiniana* occurred near the water discharge vector. Their relations with the water discharge suggest that these species might have been drifted from the bottom of the feeding river. The highest water discharge occurred in the spring when these species were dominant in the reservoir. The flow rate in rivers is probably the most effective factor controlling the population density of diatoms in the inlets of lakes. Baykal *et al.* (2011) found out that *Stephanodiscus* species were abundant in Melen River, Turkey. They state that *Stephanodiscus* species are well adapted to turbulent and turbid river systems with high nutrient concentrations.

Bere and Tundisi (2011) observed high abundance of benthic *C. meneghiniana* in the eutrophic Monjolinho River, Brazil. They state that certain benthic diatoms are associated with eutrophication and may be used as indicator species of eutrophication in running waters. Although the phytoplankton of the feeding river was not explored during the study, the high nutrient concentration of the Simav Stream might have favored high abundance diatoms in the system (Gunduzet *et al.*, 2010).

*G. subconstricta* (Chlorophyta) and *Mougeotiasp.* (Streptophyta) were dominant during the fall. In the CCA diagram, these species occurred near the water temperature and  $NH_4$ -N vectors. They had significant negative correlations with Secchi disk transparency and significant positive correlations with  $PO_4$ . The dominance of these filamentous green algae in October seems to be related to the fall overturn in the Çaygören Reservoir. In the fall, nutrients are increased and the transparency is decreased due to the overturn in the reservoir.

*C. pyrenoidifera* (Cryptophyta) and *P. nannoplanctica* (Cryptophyta) dominated phytoplankton during the winter time. These species had significant negative correlations with water temperature. Low water temperatures and low light availability may have acted as selecting factors for

this group during the winter since Cryptophytes are adapted to a low light intensity (Barone and Naselli-Flores, 2003). Various factors may regulate Cryptophyta seasonality in lakes, but it seems that the key factor in the success of Cryptophyta species is their low light requirement. Low Secchi disk transparency, during the winter in the Çaygören Reservoir, might have favored the success of this group during the winter.

The maximum phytoplankton abundance was measured in the summer because of sufficient light and high water temperature and the minimum abundance was measured in the winter because of insufficient light and low water temperature. In temperate lakes, low winter irradiance and water temperature preclude the development of high phytoplankton density in the winter time (Peeters *et al.*, 2007). The high phytoplankton densities in the summer were probably attributed to the dominance of cyanobacteria (over 80%). The high abundance of cyanobacteria during warm seasons in the Çaygören Reservoir can be attributed to the increased water temperature.

Although the maximum phytoplankton density was measured in the summer, the maximum biomass was measured in the spring. This was probably due to the dominance of cyanobacteria in the summer. Cyanobacteria have a smaller cell size than most of the other phytoplankton groups (Ciottiet *al.*, 2002). Therefore, high abundance of cyanobacteria might have not resulted in a high phytoplankton biomass in the Çaygören Reservoir.

## 5. Conclusions

The present study revealed that the important factors affecting the density, biomass and dominance of the phytoplankton in a temperate eutrophic reservoir were water temperature, underwater light (transparency), water discharge and the relative concentrations of nutrients. High density of cyanobacteria does always not warrant high biomass in the eutrophic freshwater systems. The flow rate of feeding rivers can significantly affect the population density of diatoms in the inlets of reservoirs. Finally, low water temperatures and low light availability may favor Cryptophyta dominance in the eutrophic temperate lakes.

## Acknowledgments

The present study was supported by Balikesir University Research Foundation (Project Number: 2007/18).

## References

A.P.H.A. 1995. **Standard methods for the examination of water and waste water**, 19th Ed. A.P.H.A. 1015 fifteenth Street, NW. Washington, DC 20005-2605.

Anagnostidis and Komarek J. 1988. Modern approach to the classification system of cyanophytes 3 Oscillatoriales. *Arch Hydrobiol (Supply, Algological Studies)*, **50**: 327-472.

Arhonditsis GB, Winder M, Brett MT, and Schindler DE. 2004. Patterns and mechanisms of phytoplankton variability in Lake Washington (USA). *Water Res*, **38**: 4013-4027.

Barone and Naselli-Flores LC. 2003. Distribution and seasonal dynamics of Cryptomonads in Sicilian water bodies. *Hydrobiologia*, **502**: 325-329.

Baykal T, Açıköz I and Yıldız K. 2011. Seasonal variations in phytoplankton composition and biomass in a small lowland river-lake system (Melen River, Turkey). *Turkish J Biol*, **35**: 485-501.

Bere and Tundisi JG. 2011. Influence of ionic strength and conductivity on benthic diatom communities in a tropical river (Monjolinho), Sao Carlos-SP, Brazil. *Hydrobiologia*, **661**: 261-276.

Bourrelly P. 1966. **Les algues d'eau douce. Tome I. Les algues vertes** [in French]. (Edn. N. Boubee and Cie). Société Nouvelle des, France.

Caroni R, Free G, Visconti A and Manca M. 2012. Phytoplankton functional traits and seston stable isotopes signature: a functional-based approach in a deep, subalpine lake, Lake Maggiore (N. Italy). *J Limnol*, **71** (1): 84-94.

Chaffin JD, Bridgeman TB, Heckathorn SA and Mishra S. 2011. Assessment of *Microcystis* growth rate potential and nutrient status across a trophic gradient in western Lake Erie. *J Great Lakes Res*, **37**: 92-100.

Ciotti AM, Lewis MR and Cullen JJ. 2002. Assessment of the Relationships between Dominant Cell Size in Natural Phytoplankton Communities and the Spectral Shape of the Absorption Coefficient. *Limnol Oceanogr*, **47**: 404-417.

Edmondson WT. 1971. **Productivity in freshwaters**. IPB Handbook No 17. Blackwell, UK.

Elliott JA. 2012. Predicting the impact of changing nutrient load and temperature on the phytoplankton of England's largest lake, Windermere. *Freshwater Biol*, **57**: 400-413.

Fathi AA, Abdelzaher HMA, Flower RJ, Ramdani M and Kraiem MM. 2001. Phytoplankton communities of North African wetland lakes: the CASSARINA Project. *Aquat Ecol*, **35**: 303-318.

Feuchtmayr H, Thackeray SJ, Jones ID, Ville M, Fletcher J, James J and Kelly J. 2012. Spring phytoplankton phenology - are patterns and drivers of change consistent among lakes in the same climatological region? *Freshwater Biol*, **57**: 331-34.

George DG. 1993. Physical and chemical scales of pattern in freshwater lakes and reservoirs. *Sci Total Environ*, **135**: 1-15.

Gunduz O, Simsek C and Hasozbek A. 2010. Arsenic pollution in the groundwater of Simav Plain, Turkey: Its impact on water quality and human health. *Water Air Soil Pollut*, **205**: 43-62.

Haande S, Rohrlack T, Semyalo RP, Brettum P, Edvardsen B, Lyche-Solheim A, Sorensen K and Larsson P. 2011. Phytoplankton dynamics and cyanobacterial dominance in Murchison Bay of Lake Victoria (Uganda) in relation to environmental conditions. *Limnologia*, **41**: 20-29.

House WA and Deniso FH. 2000. Factors influencing the measurement of equilibrium phosphate concentration in river sediments. *Water Res*, **34**: 1187-1200.

- Huber Pestalozzi G. 1941. **Das Phytoplankton des Süßwassers.** (Die Binnengewässer, B and XVI). Teil 2. (I) Chrysophyceen, Farblose Flagellaten Heterokonten [in German]. E. Schweizerbart'sche Verlag-sbuchhandlung, Stuttgart, Germany.
- Huber Pestalozzi G. 1950. **Das Phytoplankton des Süßwassers, Cryptophyceen, Chloromonadien, Peridiniien.** 3.Teil. (Ed. by A. Thienemann) [in German]. E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, Germany.
- Huber Pestalozzi G. 1969. **Das Phytoplankton des Süßwassers, systematik und biologie, Part 4: Euglenophyceen** [in German]. E. Schweizerbart'sche Verlag-sbuchhandlung, Stuttgart, Germany.
- Huber Pestalozzi G. 1972. **Das Phytoplankton des Süßwassers, systematik und biologie, Part 6, Fott. B. Chlorophyceae** (Grünalgen), Ordnung Tetrasporales [in German]. E. Schweizerbart'sche Verlag-sbuchhandlung, Stuttgart, Germany.
- Huber Pestalozzi G. 1982. **Das phytoplankton des süßwassers systematik und biologie, 8.Teil, 1. Halffe Conjugatophyceae Zygnematales und Desmidiaceae (excl. Zygnemataceae)** [in German]. E. Schweizerbart'sche Verlag-sbuchhandlung, Stuttgart, Germany.
- Huber Pestalozzi G. 1983. **Das phytoplankton des süßwasserssystematik und biologie, 7. Teil, 1. Halffe Chlorophyceae (Grünalgen) Ordnung: Chlorococcales** [in German]. E. Schweizerbart'sche Verlag-sbuchhandlung, Stuttgart, Germany.
- Hutchinson GE. 1967. **A treatise on limnology (vol. II): Introduction to lake biology and the limnoplankton.** John Wiley and sons. Inc., England.
- Jeppesen E, Kronvang B, Olesen JE, Audet J, Søndergaard M, Hoffmann CC, Andersen HE, Lauridsen TL, Liboriussen L, Larsen SE, Beklioglu M, Meerhoff M, Özen A and Özkan K. 2011. Climate change effects on nitrogen loading from cultivated catchments in Europe: implications for nitrogen retention, ecological state of lakes and adaptation. *Hydrobiologia*, **663**:1-21.
- John DM, Whitton BA and Brook AJ. 2003. **The Freshwater Algal Flora of the British Isles: An Identification Guide to Freshwater and Terrestrial Algae.** The Natural History Museum and the British Phycological Society. Cambridge University Press, England.
- Jones-Lee A and Lee FG. 2005. **Eutrophication (Excessive Fertilization)** In: Water Encyclopedia: Surface and Agricultural Water. Wiley, USA.
- Komarek J and Anagnostidis K. 1986. Modern approach to the classification system of cyanophytes 2. Chroococcales. *Arch Hydrobiol*, **43**: 157-226.
- Komarek J and Anagnostidis K. 1989. Modern approach to the classification system of cyanophytes. 4. Nostocales. *Arch. Hydrobiol. (Beiheft Ergebnisse der Limnologie)*, **56**: 247-345.
- Komarek J and Anagnostidis K. 1999. **Cyanoprokaryota. 1. Teil: Chroococcales.** E. Schweizerbart'sche Verlag-sbuchhandlung, Stuttgart, Germany.
- Komarek J and Anagnostidis K. 2008. **Cyanoprokaryota, 2. Teil/Part 2: Oscillatoriales, Süßwasser Flora von Mitteleuropa (Freshwater Flora of Central Europe).** E. Schweizerbart'sche Verlag-sbuchhandlung, Stuttgart, Germany.
- Krammer K and Lange-Bertalot H. 1991. **Bacillariophyceae. 4. Teil: Achnantheaceae. Kritische Ergänzungen zu Navicula (Lineolatae) und Gomphonema. Gesamtliteraturverzeichnis. Teil 1-4.** In: Ettl, H., Gerloff, J., Heynig, H. and Mollenhauer, D. (eds): **Süßwasserflora von Mitteleuropa. 2 (4).** E. Schweizerbart'sche Verlag-sbuchhandlung, Stuttgart, Germany.
- Krammer K and Lange-Bertalot H. 1999. **Bacillariophyceae. 1. Teil: Naviculaceae.** Durchgesehener Nachdruck der 1. Auflage. Spektrum Akademischer Verlag, Heidelberg, Germany.
- Kruk C, Mazzeo N, Lacerot G and Reynolds CS. 2002. Classification schemes for phytoplankton: a local validation of a functional approach to the analysis of species temporal replacement. *J Plankton Res*, **24**: 901-912.
- LaBounty FJ. 2008. Secchi transparency of Boulder Basin, Lake Mead, Arizona-Nevada: 1990-2007. *Lake Reserv Manage*, **24**: 207-218.
- Liikanen A and Martikainen PJ. 2003. Effect of ammonium and oxygen on methane and nitrous oxide fluxes across sediment-water interface in a eutrophic lake. *Chemosphere*, **52**: 1287-1293.
- Munawar M and Munawar IF. 1986. The seasonality of phytoplankton in the North American Great Lakes, a comparative synthesis. *Hydrobiologia*, **138**: 85-115.
- Niu Y, Shen H, Chen J, Xie P, Yang X, Tao M, Ma Z and Qi M. 2011. Phytoplankton community succession shaping bacterioplankton community composition in Lake Taihu, China. *Water Res.*, **45**: 4169-4182.
- Padisak J, Crossetti LO and Naselli-Flores L. 2009. Use and misuse in the application of the phytoplankton functional classification: a critical review with updates. *Hydrobiologia*, **621**: 1-19.
- Parinet B, Lhote A and Legube B. 2004. Principal component analysis: an appropriate tool for water quality evaluation and management-application to a tropical lake system. *Ecol Model*, **178**: 295-311.
- Peeters F, Straile D, Lorke A and Ollinger D. 2007. Turbulent Mixing and Phytoplankton Spring Bloom Development in a Deep Lake. *Limnol Oceanogr*, **52**: 286-298.
- Pérez G, Queimaliños C, Balseiro E and Modenutti B. 2007. Phytoplankton absorption spectra along the water column in deep North Patagonian Andean lakes (Argentina). *Limnologia*, **37**: 3-16.
- Reynolds CS, Huszar V, Kruk C, Naselli-Flores L and Melo S. 2002. Towards a functional classification of the freshwater phytoplankton. *J Plankton Res*, **24**: 417-428.
- Round FE, Crawford RM and Mann DG. 1990. **The Diatoms: Morphology and biology of the genera.** Cambridge University Press, England.
- SAS Institute. 2003. **SAS/STAT Users Guide** (Version 9.1). SAS Institute Inc., USA.
- Scheffer M and van Nes EH. 2007. Shallow lakes theory revisited. Various alternative regimes driven by climate, nutrients, depth and lake size. *Hydrobiologia*, **584**: 455-466.