

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/257827010>

Commutator Subgroups of the Extended Hecke Groups $\bar{H}(\lambda, q)$

Article in *Czechoslovak Mathematical Journal* · March 2004

DOI: 10.1023/B:CMAJ.0000027265.81403.8d

CITATIONS

18

READS

56

3 authors:



Recep Sahin

Balikesir University

38 PUBLICATIONS 143 CITATIONS

[SEE PROFILE](#)



Osman Bizim

Uludag University

21 PUBLICATIONS 77 CITATIONS

[SEE PROFILE](#)



Ismail naci Cangul

Uludag University

171 PUBLICATIONS 1,576 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



co pi energy [View project](#)



Studies in Some Algebraic and Topological Properties of Graphs [View project](#)

COMMUTATOR SUBGROUPS OF THE EXTENDED
HECKE GROUPS $\overline{H}(\lambda_q)$

R. SAHIN, Balikesir, O. BIZIM and I. N. CANGUL, Bursa

(Received July 9, 2001)

Abstract. Hecke groups $H(\lambda_q)$ are the discrete subgroups of $\text{PSL}(2, \mathbb{R})$ generated by $S(z) = -(z + \lambda_q)^{-1}$ and $T(z) = -1/z$. The commutator subgroup of $H(\lambda_q)$, denoted by $H'(\lambda_q)$, is studied in [2]. It was shown that $H'(\lambda_q)$ is a free group of rank $q - 1$.

Here the extended Hecke groups $\overline{H}(\lambda_q)$, obtained by adjoining $R_1(z) = 1/\bar{z}$ to the generators of $H(\lambda_q)$, are considered. The commutator subgroup of $\overline{H}(\lambda_q)$ is shown to be a free product of two finite cyclic groups. Also it is interesting to note that while in the $H(\lambda_q)$ case, the index of $H'(\lambda_q)$ is changed by q , in the case of $\overline{H}(\lambda_q)$, this number is either 4 for q odd or 8 for q even.

Keywords: Hecke group, extended Hecke group, commutator subgroup

MSC 2000: 11F06, 20H05, 20H10

1. INTRODUCTION

In [4], Erich Hecke introduced the groups $H(\lambda)$ generated by two linear fractional transformations

$$T(z) = -\frac{1}{z} \quad \text{and} \quad U(z) = z + \lambda,$$

where λ is a fixed positive real number. T and U have matrix representations

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix},$$

respectively. (In this work we identify each matrix A with $-A$, so that they each represent the same transformation). Let $S = T.U$, i.e.

$$S(z) = -\frac{1}{z + \lambda}.$$

E. Hecke showed that $H(\lambda)$ is Fuchsian if and only if $\lambda = \lambda_q = 2 \cos \frac{\pi}{q}$, where q is an integer $q \geq 3$ or $\lambda > 2$ is real. In these two cases $H(\lambda)$ is called a Hecke group. We consider the former case. Then the Hecke group $H(\lambda)$ is the discrete subgroup of $\text{PSL}(2, \mathbb{R})$ generated by S and U , where

$$U(z) = z + \lambda_q$$

and it has a presentation $H(\lambda) = \langle T, S \mid T^2 = S^q = I \rangle$.

The most important and studied Hecke group is the modular group $H(\lambda_3)$. In this case $\lambda_3 = 2 \cos \frac{\pi}{3} = 1$, i.e. all coefficients of the elements of $H(\lambda_3)$ are rational integers. In the literature, the symbols Γ and $\Gamma(1)$ are used to denote the modular group. In this paper we shall use $H(\lambda_3)$ for this purpose. The next two most important Hecke groups are those for $q = 4$ and $q = 6$, in which cases $\lambda_q = \sqrt{2}$ and $\sqrt{3}$, respectively.

The extended modular group $\overline{H}(\lambda_3)$ has a presentation

$$\overline{H}(\lambda_3) = \langle R_1, R_2, R_3 \mid R_1^2 = R_2^2 = R_3^2 = (R_1 R_2)^3 = (R_3 R_1)^2 = I \rangle$$

where

$$R_1(z) = \frac{1}{\bar{z}}, \quad R_2(z) = \frac{-1}{\bar{z} + 1}, \quad R_3(z) = -\bar{z}.$$

The modular group is a subgroup of index 2 in $\overline{H}(\lambda_3)$ (see [3]). It has a presentation

$$H(\lambda_3) = \langle T, S \mid T^2 = S^3 = I \rangle \cong C_2 * C_3,$$

where

$$T = R_3 R_1 = R_1 R_3, \quad S = R_1 R_2.$$

Putting $R = R_1$, we have

$$\overline{H}(\lambda_3) = \langle T, S, R \mid T^2 = S^3 = R^2 = I, RT = TR, RS = S^{-1}R \rangle.$$

Similarly the extended Hecke group $\overline{H}(\lambda_q)$ has a presentation

$$\overline{H}(\lambda_q) = \langle T, S, R \mid T^2 = S^q = R^2 = I, RT = TR, RS = S^{-1}R \rangle$$

and Hecke group $H(\lambda_q)$ is a subgroup of index 2 in $\overline{H}(\lambda_q)$.

The commutator subgroup of G is denoted by G' and defined by

$$\langle [g, h] \mid g, h \in G \rangle$$

where $[g, h] = ghg^{-1}h^{-1}$. Since G' is a normal subgroup of G , we can form the factor-group G/G' which is the largest abelian quotient group of G .

In this work we obtain some results concerning commutator subgroups of the extended Hecke group $\overline{H}(\lambda_q)$.

2. COMMUTATOR SUBGROUPS OF THE EXTENDED HECKE GROUP $\overline{H}(\lambda_q)$

The commutator subgroup of the Hecke group $H(\lambda_q)$ is denoted by $H'(\lambda_q)$. We have

$$T^2 = S^q = I, \quad TS = ST$$

in $H(\lambda_q)/H'(\lambda_q)$. So one can find

$$H(\lambda_q)/H'(\lambda_q) \cong C_2 \times C_q$$

and hence it is isomorphic to C_{2q} if q is odd. Therefore

$$|H(\lambda_q) : H'(\lambda_q)| = 2q.$$

If q is even, $(TS)^q = 1$ while if q is odd, $(TS)^{2q} = 1$. In particular, $H'(\lambda_q)$ is a free group of rank $q - 1$ (see [1]).

By [5], the Reidemeister-Schreier method gives the generators of $H'(\lambda_q)$ as

$$a_1 = TSTS^{q-1}, \quad a_2 = TS^2TS^{q-2}, \quad \dots, \quad a_{q-1} = TS^{q-1}TS.$$

Similarly for the extended Hecke group $\overline{H}(\lambda_q)$ we have

$$T^2 = S^q = R^2 = I, \quad RT = TR, \quad RS = S^{-1}R, \quad RS = SR, \quad TS = ST$$

in $\overline{H}(\lambda_q)/\overline{H}'(\lambda_q)$.

Theorem 1. *Let q be odd, then*

- (i) $\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) \cong V_4 \cong C_2 \times C_2$
- (ii) $\overline{H}'(\lambda_q) = \langle S, TST \mid S^q = (TST)^q = I \rangle \cong C_q * C_q$.

Proof. (i) Since the extended Hecke group $\overline{H}(\lambda_q)$ has a presentation

$$\overline{H}(\lambda_q) = \langle T, S, R \mid T^2 = S^q = R^2 = I, RT = TR, RS = S^{-1}R \rangle$$

and

$$\begin{aligned} \overline{H}(\lambda_q)/\overline{H}'(\lambda_q) = \langle T, S, R \mid T^2 = S^q = R^2 = I, RT = TR, RS = S^{-1}R, \\ RS = SR, TS = ST \rangle \end{aligned}$$

one has $RS = S^{-1}R$ and $RS = SR$, and thus

$$S^{q-2} = S^q = S^2 = I.$$

This shows that $S = I$, as q is odd. Thus

$$\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) = \langle T, R \mid T^2 = R^2 = (TR)^2 = I \rangle$$

and finally

$$\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) \cong V_4 \cong C_2 \times C_2.$$

(ii) Now we determine the set of generators for $\overline{H}'(\lambda_q)$. We choose a Schreier transversal for $\overline{H}'(\lambda_q)$ as

$$I, T, R, TR.$$

According to the Reidemeister-Schreier method, we can form all possible products

$$\begin{array}{lll} I \cdot T \cdot (T)^{-1} = I, & I \cdot S \cdot (I)^{-1} = S, & I \cdot R \cdot (R)^{-1} = I, \\ T \cdot T \cdot (I)^{-1} = I, & T \cdot S \cdot (T)^{-1} = TST, & T \cdot R \cdot (TR)^{-1} = I, \\ R \cdot T \cdot (TR)^{-1} = RTRT, & R \cdot S \cdot (R)^{-1} = RSR, & R \cdot R \cdot (I)^{-1} = I, \\ TR \cdot T \cdot (R)^{-1} = TRTR, & TR \cdot S \cdot (TR)^{-1} = TRSRT, & TR \cdot R \cdot (T)^{-1} = I. \end{array}$$

Since

$$\begin{array}{l} RTRT = I, \\ TRTR = I, \\ RSR = S^{-1}, \\ TRSRT = TS^{-1}T = (TST)^{-1}, \end{array}$$

the generators are S and TST . Thus $\overline{H}'(\lambda_q)$ has a presentation

$$\overline{H}'(\lambda_q) = \langle S, TST \mid S^q = (TST)^q = I \rangle \cong C_q * C_q.$$

□

Theorem 2. *Let q be even, then*

- (i) $\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) \cong C_2 \times C_2 \times C_2$
- (ii) $\overline{H}'(\lambda_q) = \langle S^2, TS^2T, TST S^{q-1} \mid (S^2)^{q/2} = (TS^2T)^{q/2} = (TST S^{q-1})^\infty = I \rangle$.

Proof. (i) If the representations of $\overline{H}(\lambda_q)$ and $\overline{H}(\lambda_q)/\overline{H}'(\lambda_q)$ are considered, we obtain $S^2 = I$ as $RS = S^{-1}R$ and $RS = SR$, $S^{q-2} = S^q = S^2 = I$ as q is odd. Therefore

$$\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) = \langle T, S, R \mid T^2 = S^2 = R^2 = (RT)^2 = (RS)^2 = (TS)^2 = I \rangle$$

and so

$$\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) \cong C_2 \times C_2 \times C_2.$$

(ii) Again we choose a Schreier transversal for $\overline{H}'(\lambda_q)$ as

$$I, T, R, S, TR, SR, TS, TSR.$$

Hence, all possible products are

$$\begin{aligned} I \cdot T \cdot (T)^{-1} &= I, & TR \cdot T \cdot (R)^{-1} &= TRTR, \\ T \cdot T \cdot (I)^{-1} &= I, & SR \cdot T \cdot (TSR)^{-1} &= SRTRS^{-1}T, \\ R \cdot T \cdot (TR)^{-1} &= RTRT, & TS \cdot T \cdot (S)^{-1} &= TSTS^{-1}, \\ S \cdot T \cdot (TS)^{-1} &= STS^{-1}T, & TSR \cdot T \cdot (SR)^{-1} &= TSRTRS^{-1}, \\ I \cdot S \cdot (S)^{-1} &= I, & TR \cdot S \cdot (TSR)^{-1} &= TRSRS^{-1}T, \\ T \cdot S \cdot (TS)^{-1} &= I, & SR \cdot S \cdot (R)^{-1} &= SRSR, \\ R \cdot S \cdot (SR)^{-1} &= RSRS^{-1}, & TS \cdot S \cdot (T)^{-1} &= TS^2T, \\ S \cdot S \cdot (I)^{-1} &= S^2, & TSR \cdot S \cdot (TR)^{-1} &= TSRSRT, \\ I \cdot R \cdot (R)^{-1} &= I, & TR \cdot R \cdot (T)^{-1} &= I, \\ T \cdot R \cdot (TR)^{-1} &= I, & SR \cdot R \cdot (S)^{-1} &= I, \\ R \cdot R \cdot (I)^{-1} &= I, & TS \cdot R \cdot (TSR)^{-1} &= I, \\ S \cdot R \cdot (SR)^{-1} &= I, & TSR \cdot R \cdot (TS)^{-1} &= I. \end{aligned}$$

Since $(STS^{-1}T)^{-1} = TSTS^{-1}$, $(TRTR)^{-1} = RTRT = I$, $(RSRS^{-1}) = (S^2)^{-1}$, $SRSR = I$, $SRTRS^{-1}T)^{-1} = TSRTRS^{-1} = TSTS^{-1}$, $TRRS^{-1}T = (TS^2T)^{-1}$, $TSRSRT = I$, the generators of $\overline{H}'(\lambda_q)$ are S^2 , TS^2T , $TSTS^{q-1}$. Thus $\overline{H}'(\lambda_q)$ has a presentation

$$\overline{H}'(\lambda_q) = \langle S^2 \rangle * \langle TS^2T \rangle * \langle TSTS^{q-1} \rangle.$$

Example 1. Let $q = 3$. Then $\overline{H}(\lambda_3)$ is the extended modular group. In this case

$$\overline{H}(\lambda_3)/\overline{H}'(\lambda_3) = \langle T, R \mid T^2 = R^2 = (TR)^2 = I \rangle$$

and a Schreier transversal is

$$I, T, R, TR.$$

Hence,

$$\begin{aligned} I \cdot T \cdot (T)^{-1} &= I, & I \cdot S \cdot (I)^{-1} &= S, & I \cdot R \cdot (R)^{-1} &= I, \\ T \cdot T \cdot (I)^{-1} &= I, & T \cdot S \cdot (T)^{-1} &= TST, & T \cdot R \cdot (TR)^{-1} &= I, \\ R \cdot T \cdot (TR)^{-1} &= RTRT, & R \cdot S \cdot (R)^{-1} &= RSR, & R \cdot R \cdot (I)^{-1} &= I, \\ TR \cdot T \cdot (R)^{-1} &= TRTR, & TR \cdot S \cdot (TR)^{-1} &= TRSRT, & TR \cdot R \cdot (T)^{-1} &= I \end{aligned}$$

and since $RTRT = I$, $TRTR = I$, $RSR = S^{-1}$, $TRSRT = TS^{-1}T = (TST)^{-1}$, the generators of $\overline{H}(\lambda_3)$ are S and TST . Thus $\overline{H}'(\lambda_3)$ has a presentation

$$\overline{H}'(\lambda_3) = \langle S, TST \mid S^3 = (TST)^3 = I \rangle \cong C_3 * C_3.$$

Notice that this result coincides with the ones given in [5] for the extended modular group.

Example 2. Let $q = 6$. Then $\overline{H}(\lambda_6)$ and $\overline{H}(\lambda_6)/\overline{H}'(\lambda_6)$ have presentations

$$\overline{H}(\lambda_6) = \langle T, S, R \mid T^2 = S^6 = R^2 = I, TR = RT, RS = S^{-1}R \rangle$$

and

$$\begin{aligned} \overline{H}(\lambda_6)/\overline{H}'(\lambda_6) = \langle T, S, R \mid T^2 = S^6 = R^2 = I, RT = TR, RS = S^{-1}R, \\ RS = SR, TS = ST \rangle. \end{aligned}$$

Since $RS = S^{-1}R$ and $RS = SR$, $S^{-1} = S^5$ and so $S^4 = S^6 = I$, $S^2 = I$. Hence

$$\overline{H}(\lambda_6)/\overline{H}'(\lambda_6) = \langle T, S, R \mid T^2 = S^2 = R^2 = (RT)^2 = (RS)^2 = (TS)^2 = I \rangle.$$

We can choose a Schreier transversal as

$$I, T, R, S, TR, SR, TS, TSR.$$

In this case all the possibilities are

$$\begin{array}{ll} I \cdot T \cdot (T)^{-1} = I, & TR \cdot T \cdot (R)^{-1} = TRTR, \\ T \cdot T \cdot (I)^{-1} = I, & SR \cdot T \cdot (TSR)^{-1} = SRTRS^5T, \\ R \cdot T \cdot (TR)^{-1} = RTRT, & TS \cdot T \cdot (S)^{-1} = TSTS^5, \\ S \cdot T \cdot (TS)^{-1} = STS^5T, & TSR \cdot T \cdot (SR)^{-1} = TSRTRS^5, \\ I \cdot S \cdot (S)^{-1} = I, & TR \cdot S \cdot (TSR)^{-1} = TRSR S^5T, \\ T \cdot S \cdot (TS)^{-1} = I, & SR \cdot S \cdot (R)^{-1} = SRSR, \\ R \cdot S \cdot (SR)^{-1} = RSRS^5, & TS \cdot S \cdot (T)^{-1} = TS^2T, \\ S \cdot S \cdot (I)^{-1} = S^2, & TSR \cdot S \cdot (TR)^{-1} = TSRSRT, \\ I \cdot R \cdot (R)^{-1} = I, & TR \cdot R \cdot (T)^{-1} = I, \\ T \cdot R \cdot (TR)^{-1} = I, & SR \cdot R \cdot (S)^{-1} = I, \\ R \cdot R \cdot (I)^{-1} = I, & TS \cdot R \cdot (TSR)^{-1} = I, \\ S \cdot R \cdot (SR)^{-1} = I, & TSR \cdot R \cdot (TS)^{-1} = I. \end{array}$$

Since $(STS^5T)^{-1} = TSTS^5$, $(TRTR)^{-1} = TRTR = I$, $(RSRS^5) = (S^2)^{-1}$, $SRSR = I$, $(SRTRS^5T)^{-1} = TSRTRS^5 = TSTS^5$, $TRRS^5T = (TS^2T)^{-1}$, $TSRSRT = I$, the generators of $\bar{H}'(\lambda_q)$ are $S^2, TS^2T, TSTS^5$. Thus $\bar{H}'(\lambda_6)$ has a presentation

$$\bar{H}'(\lambda_6) = \langle S^2 \rangle * \langle TS^2T \rangle * \langle TSTS^5 \rangle.$$

References

- [1] *R. B. J. T. Allenby*: Rings, Fields and Groups. Second Edition. Edward Arnold, London-New York-Melbourne-Auckland, 1991.
- [2] *I. N. Cangül and D. Singerman*: Normal subgroups of Hecke groups and regular maps. Math. Proc. Camb. Phil. Soc. *123* (1998), 59–74.
- [3] *H. S. M. Coxeter and W. O. J. Moser*: Generators and Relations for Discrete Groups. Springer, Berlin, 1957.
- [4] *E. Hecke*: Über die Bestimmung Dirichletscher Reihen durch ihre Funktionalgleichungen. Math. Ann. *112* (1936), 664–699.
- [5] *D. L. Johnson*: Topics in the Theory of Group Presentations. L.M.S. Lecture Note Series 42. Cambridge Univ. Press, Cambridge, 1980.
- [6] *G. A. Jones and J. S. Thornton*: Automorphisms and congruence subgroups of the extended modular group. J. London Math. Soc. *34* (1986), 26–40.

Authors' addresses: R. Sahin, Balikesir Universitesi Fen-Edebiyat Fakültesi Matematik Bölümü 10100, Balikesir, Turkey, e-mail: rsahin@balikesir.edu.tr; O. Bizim, I. N. Cangül, Uludag Universitesi Fen-Edebiyat Fakültesi Matematik Bölümü Görükle 16059, Bursa, Turkey, e-mails: obizim@uludag.edu.tr, cangul@uludag.edu.tr.