



On the efficiency of some p -groups

Firat Ateş

Abstract

Let p be a prime number. In this paper, we work on the efficiency of the p -groups G_1 and G_2 defined by the presentations,

$$\mathcal{P}_{G_1} = \langle a, b, c ; ab = bac, bc = cb, ac = ca, a^{p^\alpha} = 1, b^{p^\beta} = 1, c^{p^\gamma} = 1 \rangle$$

where $\alpha \geq \beta > \gamma \geq 1$ and

$$\mathcal{P}_{G_2} = \langle a, b ; ab = ba^{1+p^{\alpha-\gamma}}, a^{p^\alpha} = 1, b^{p^\beta} = 1 \rangle$$

where $\alpha \geq 2\gamma$, $\beta > \gamma \geq 1$ and $\alpha + \beta > 3$. For example, if we let $p = 2$, then by [1], the groups defined by these presentations becomes 2-groups. It is known that these groups play an important role in the theory of groups of nilpotency class 2.

1 Introduction

Let G be a finitely presented group with a presentation

$$\mathcal{P} = \langle \mathbf{x} ; \mathbf{r} \rangle. \quad (1)$$

Then the deficiency of this presentation is defined by $|\mathbf{r}| - |\mathbf{x}|$, and is denoted by $def(\mathcal{P})$. Moreover, the group deficiency of a finitely presented group G is given by

$$def_G(G) = \min\{def(\mathcal{P}) : \mathcal{P} \text{ is a finite group presentation for } G\}.$$

Key Words: Efficiency, pictures, p -groups
2010 Mathematics Subject Classification: Primary 20E22, 20J05; Secondary 20F05, 57M05.

Received: February, 2014.

Revised: May, 2014.

Accepted: May, 2014.

One can apply similar definitions for the semigroup deficiency of a finitely presented semigroup S , $def_S(S)$. Let us consider the second integral homology $H_2(G)$ of a finite group G . It is well known that the group G (or semigroup S) is efficient as a group (or as a semigroup), if we have $def_G(G) = rank(H_2(G))$ (or $def_S(S) = rank(H_2(S^1))$) where S^1 is obtained from S by adjoining an identity). We can refer to the reader [2, 3, 8, 9, 10] for more details.

One of the most effective way to show efficiency for the group G is to use *spherical pictures* ([7, 18]) over \mathcal{P} . These geometric configurations are the representative elements of the second homotopy group $\pi_2(\mathcal{P})$ of \mathcal{P} which is a left $\mathbb{Z}G$ -module. They are denoted by \mathbb{P} .

Suppose \mathbf{Y} is a collection of spherical pictures over \mathbb{P} . Then, by [18], one can define the additional operation on spherical pictures. Allowing this additional operation leads to the notion of *equivalence (rel \mathbf{Y}) of spherical pictures*. Then, again in [18], Pride proved that *the elements $\langle \mathbb{P} \rangle$ ($\mathbb{P} \in \mathbf{Y}$) generate $\pi_2(\mathcal{P})$ as a module if and only if every spherical picture is equivalent (rel \mathbf{Y}) to the empty picture*. Therefore one can easily say that if the elements $\langle \mathbb{P} \rangle$ ($\mathbb{P} \in \mathbf{Y}$) generate $\pi_2(\mathcal{P})$, then \mathbf{Y} generates $\pi_2(\mathcal{P})$.

For any picture \mathbb{P} over \mathcal{P} and for any $R \in \mathbf{r}$, the *exponent sum* of R in \mathbb{P} , denoted by $exp_R(\mathbb{P})$, is the number of discs of \mathbb{P} labeled by R minus the number of discs labeled by R^{-1} . We remark that if any two pictures \mathbb{P}_1 and \mathbb{P}_2 are equivalent then $exp_R(\mathbb{P}_1) = exp_R(\mathbb{P}_2)$, for all $R \in \mathbf{r}$. Let n be a non-negative integer. Then \mathcal{P} is said to be *n -Cockcroft* if $exp_R(\mathbb{P}) \equiv 0 \pmod{n}$, (where congruence $\pmod{0}$ is taken to be equality) for all $R \in \mathbf{r}$ and for all spherical pictures \mathbb{P} over \mathcal{P} . Then a group G is said to be *n -Cockcroft* if it admits an n -Cockcroft presentation. To verify that the n -Cockcroft property holds, it is enough to check for pictures $\mathbb{P} \in \mathbf{Y}$, where \mathbf{Y} is a set of generating pictures. The case $n = 0$ is just called Cockcroft. One can refer [11], [13], [14], [15] and [17] for the Cockcroft property and [9], [17] for the n -Cockcroft property.

The subject *efficiency*, for the presentation \mathcal{P} as in (1) and so for the group G , is related to the q -Cockcroft property (see Theorem 1.1 below). We can refer, for example, [4] and [10] for the definition and applications of efficiency. We then have the following result.

Theorem 1.1 ([12, 17]). *Let \mathcal{P} be as in (1). Then \mathcal{P} is efficient if and only if it is q -Cockcroft for some prime q .*

2 Main results

In [6], Bacon and Kappe worked on two-generator p -groups of nilpotency class 2 where $p \neq 2$. Also, in [16], Kappe, Sarmin and Visscher worked on

two-generator 2-groups of nilpotency class 2. Also let us consider the following semigroups defined by the presentations:

$$\langle a, b, c ; ab = bac, bc = cb, ac = ca, a^{p^\alpha+1} = a, b^{p^\beta+1} = b, c^{p^\gamma+1} = c \rangle \quad (2)$$

where $\alpha \geq \beta \geq \gamma \geq 1$ and

$$\langle a, b ; ab = ba^{1+p^{\alpha-\gamma}}, a^{p^\alpha+1} = a, b^{p^\beta+1} = b \rangle \quad (3)$$

where $\alpha \geq 2\gamma$, $\beta \geq \gamma \geq 1$ and $\alpha + \beta > 3$. In [1], the authors showed that the semigroups defined by the presentations (2) and (3) have the orders

$$p^{\alpha+\beta+\gamma} + p^\alpha + p^\beta + p^\gamma + p^{\alpha+\beta} + p^{\beta+\gamma} + p^{\alpha+\gamma} \quad \text{and} \quad p^{\alpha+\beta} + p^\alpha + p^\beta,$$

respectively.

Now let us again think the following presentations for the groups G_1 and G_2 which are given in abstract

$$\mathcal{P}_{G_1} = \langle a, b, c ; ab = bac, bc = cb, ac = ca, a^{p^\alpha} = 1, b^{p^\beta} = 1, c^{p^\gamma} = 1 \rangle \quad (4)$$

where $\alpha \geq \beta \geq \gamma \geq 1$ and

$$\mathcal{P}_{G_2} = \langle a, b ; ab = ba^{1+p^{\alpha-\gamma}}, a^{p^\alpha} = 1, b^{p^\beta} = 1 \rangle \quad (5)$$

where $\alpha \geq 2\gamma$, $\beta \geq \gamma \geq 1$ and $\alpha + \beta > 3$. In [1], the authors showed that the groups defined by the presentations (4) and (5) have the orders

$$p^{\alpha+\beta+\gamma} \quad \text{and} \quad p^{\alpha+\beta}.$$

In this paper, our aim is to study on the efficiency of the groups G_1 and G_2 presented by (4) and (5), by using the works given [2, 3, 5, 8, 9, 10].

Therefore we can give the main results of this paper as follows.

Theorem 2.1. *For every prime number p and integers α , β and γ with $\alpha \geq \beta > \gamma \geq 1$, the group G_1 presented by (4) is efficient.*

Theorem 2.2. *For every prime number p and integers α , β and γ with $\alpha \geq 2\gamma$, $\beta > \gamma \geq 1$ and $\alpha + \beta > 3$, the group G_2 presented by (5) is efficient.*

3 Proof of the main results

3.1 Proof of Theorem 2.1

Consider the group G_1 . Since we have the following relations $ab = bac, bc = cb, ac = ca, a^{p^\alpha} = 1, b^{p^\beta} = 1, c^{p^\gamma} = 1$, we have to think about the following




Figure 1

overlapping word pairs ab^{p^β} , $a^{p^\alpha}b$, ac^{p^γ} , bc^{p^γ} , $a^{p^\alpha}c$ and $b^{p^\beta}c$ for defining the elements of $\pi_2(\mathcal{P}_{G_1})$. It is known that spherical pictures which are obtained from the resolutions of these pairs give the elements of $\pi_2(\mathcal{P}_{G_1})$ by [5].

Now, let us consider the pairs ab^{p^β} and $a^{p^\alpha}b$. Then by using the relations of the group G_1 , the resolutions for these pairs can be given as pictures \mathbf{P}_1 and \mathbf{P}_2 , respectively in Figure 1.

Now, let us also consider the discs in the pictures \mathbf{P}_1 and \mathbf{P}_2 . To prove this theorem, we need to count the exponent sums of the discs in these pictures. So let us calculate the number of S_1 -discs, S_2 -discs, S_3 -discs, S_4 -discs, S_5 -discs and S_6 -discs in \mathbf{P}_1 , \mathbf{P}_2 where $S_1 : b^{p^\beta} = 1$, $S_2 : c^{p^\gamma} = 1$, $S_3 : ab = bac$, $S_4 : a^{p^\alpha} = 1$, $S_5 : bc = cb$ and $S_6 : ac = ca$. At this point, it can be seen that

$$\begin{aligned} \text{exp}_{S_1}(\mathbf{P}_1) &= 1 - 1 = 0, & \text{exp}_{S_2}(\mathbf{P}_1) &= p^{\beta-\gamma}, \\ \text{exp}_{S_2}(\mathbf{P}_2) &= p^{\alpha-\gamma}, & \text{exp}_{S_3}(\mathbf{P}_1) &= p^\beta, \\ \text{exp}_{S_3}(\mathbf{P}_2) &= p^\alpha, & \text{exp}_{S_4}(\mathbf{P}_2) &= 1 - 1 = 0, \\ \text{exp}_{S_5}(\mathbf{P}_1) &= 1 + 2 + 3 + \dots + (p^\beta - 1) = \frac{(p^\beta - 1)p^\beta}{2}, \\ \text{exp}_{S_6}(\mathbf{P}_2) &= 1 + 2 + 3 + \dots + (p^\alpha - 1) = \frac{(p^\alpha - 1)p^\alpha}{2} \end{aligned}$$




Figure 2

and to q -Cockcroft property be hold for some prime q , we need to have

$$\begin{aligned}
 \exp_{S_2}(\mathbf{P}_1) \equiv 0 \pmod{q} &\Leftrightarrow p^{\beta-\gamma} \equiv 0 \pmod{q}, \\
 \exp_{S_2}(\mathbf{P}_2) \equiv 0 \pmod{q} &\Leftrightarrow p^{\alpha-\gamma} \equiv 0 \pmod{q}, \\
 \exp_{S_3}(\mathbf{P}_1) \equiv 0 \pmod{q} &\Leftrightarrow p^\beta \equiv 0 \pmod{q}, \\
 \exp_{S_3}(\mathbf{P}_2) \equiv 0 \pmod{q} &\Leftrightarrow p^\alpha \equiv 0 \pmod{q}, \\
 \exp_{S_5}(\mathbf{P}_1) \equiv 0 \pmod{q} &\Leftrightarrow \frac{(p^\beta - 1)p^\beta}{2} \equiv 0 \pmod{q}, \\
 \exp_{S_6}(\mathbf{P}_2) \equiv 0 \pmod{q} &\Leftrightarrow \frac{(p^\alpha - 1)p^\alpha}{2} \equiv 0 \pmod{q}.
 \end{aligned}$$

Now, let us consider the pairs ac^{p^γ} and bc^{p^γ} . Then by using the relations S_2 , S_5 and S_6 , the resolutions for these pairs can be given as pictures \mathbf{P}_3 and \mathbf{P}_4 , respectively in Figure 2.

Similarly, as in the above, we need to count the exponent sums of the discs in these pictures. Therefore let us give the number of S_2 -discs, S_5 -discs and S_6 -discs in \mathbf{P}_3 , \mathbf{P}_4 as follows;

$$\begin{aligned}
 \exp_{S_2}(\mathbf{P}_3) &= 1 - 1 = 0, & \exp_{S_2}(\mathbf{P}_4) &= 1 - 1 = 0, \\
 \exp_{S_5}(\mathbf{P}_4) &= p^\gamma, & \exp_{S_6}(\mathbf{P}_3) &= p^\gamma.
 \end{aligned}$$

So in order to give q -Cockcroft property for some prime q , we need to have

$$\exp_{S_5}(\mathbf{P}_4) = \exp_{S_6}(\mathbf{P}_3) \equiv 0 \pmod{q} \Leftrightarrow p^\gamma \equiv 0 \pmod{q}.$$

Similarly, let us consider the pairs $a^{p^\alpha}c$ and $b^{p^\beta}c$. Then by using the relations S_1 , S_4 , S_5 and S_6 , the resolutions for these pairs can be given as pictures \mathbf{P}_5 and \mathbf{P}_6 , respectively in Figure 3.




Figure 3

Here one can give the exponent sums of the discs in these pictures as follows;

$$\begin{aligned} \text{exp}_{S_1}(\mathbf{P}_6) &= 1 - 1 = 0, & \text{exp}_{S_4}(\mathbf{P}_5) &= 1 - 1 = 0, \\ \text{exp}_{S_5}(\mathbf{P}_6) &= p^\beta, & \text{exp}_{S_6}(\mathbf{P}_5) &= p^\alpha. \end{aligned}$$

Thus in order to give q -Cockcroft property for some prime q , we have

$$\begin{aligned} \text{exp}_{S_5}(\mathbf{P}_6) \equiv 0 \pmod{q} &\Leftrightarrow p^\beta \equiv 0 \pmod{q}, \\ \text{exp}_{S_6}(\mathbf{P}_5) \equiv 0 \pmod{q} &\Leftrightarrow p^\alpha \equiv 0 \pmod{q}. \end{aligned}$$




Figure 4

Also let us consider the pictures in Figure 4. Here we have

$$\text{exp}_{S_1}(\mathbf{C}_1) = 1 - 1 = 0, \quad \text{exp}_{S_2}(\mathbf{C}_2) = 1 - 1 = 0, \quad \text{exp}_{S_4}(\mathbf{C}_3) = 1 - 1 = 0.$$

Finally, we can see that $\pi_2(\mathcal{P}_{G_1})$ consists of the pictures $\mathbf{P}_1, \mathbf{P}_2, \mathbf{P}_3, \mathbf{P}_4, \mathbf{P}_5, \mathbf{P}_6, \mathbf{C}_1, \mathbf{C}_2$ and \mathbf{C}_3 . Thus in order to get q -Cockcroft property, we must calculate the exponent sums of the discs in these pictures. Then, by using the

above arguments, for getting q -Cockcroft property for some prime q , we must have

$$\begin{aligned} p^{\beta-\gamma} &\equiv 0 \pmod{q}, & p^{\alpha-\gamma} &\equiv 0 \pmod{q}, \\ p^\beta &\equiv 0 \pmod{q}, & p^\alpha &\equiv 0 \pmod{q}, & p^\gamma &\equiv 0 \pmod{q} \\ \frac{(p^\beta-1)p^\beta}{2} &\equiv 0 \pmod{q}, & \frac{(p^\alpha-1)p^\alpha}{2} &\equiv 0 \pmod{q}. \end{aligned}$$

Then by Theorem 1.1 we may say that the group G_1 is efficient if and only if it is q -Cockcroft for some prime q . At this point, since we have $\alpha \geq \beta > \gamma \geq 1$, then we choose $p = q$. This gives that the group G_1 presented by (4) is q -Cockcroft. This says that G_1 is efficient.

Remark 3.1. *We realised that we choose $\alpha \geq \beta > \gamma \geq 1$. If we choose $\alpha \geq \beta \geq \gamma \geq 1$, then we may have $\beta = \gamma$ or $\alpha = \gamma$. This gives that $p^{\beta-\gamma} = p^0 = 1$ is not equivalent to 0 by the modulo q or $p^{\alpha-\gamma} = p^0 = 1$ is not equivalent to 0 by the modulo q , for some prime q . Also, for $p = 2$, if we choose $\alpha \geq \beta \geq \gamma \geq 1$, then we may have $\beta = 1$ or $\alpha = 1$. This says that $\frac{(p^\beta-1)p^\beta}{2} = 1$ is not equivalent to 0 by the modulo q or $\frac{(p^\alpha-1)p^\alpha}{2} = 1$ is not equivalent to 0 by the modulo q , for some prime q .*

Remark 3.2. *In [3, 8], it was shown that for a finitely presented group G with non-negative deficiency we have $\text{def}_S(G) = \text{def}_G(G)$. This says that a group G with non-negative deficiency is efficient as a group if and only if G is efficient as a semigroup. Therefore, since the group G_1 presented by (4) has non-negative deficiency and it is efficient as a group, then it is also efficient as a semigroup. Hence we get that the semigroup related to the certain group presentation (2) is also efficient.*

3.2 Proof of Theorem 2.2

Let us consider the group G_2 . Here we have the following relations $a^{p^\alpha} = 1$, $b^{p^\beta} = 1$ and $ab = ba^{1+p^{\alpha-\gamma}}$. Thus we cocern about the following overlapping word pairs ab^{p^β} and $a^{p^\alpha}b$ for defining the elements of $\pi_2(\mathcal{P}_{G_2})$.

Now, let us consider the pairs ab^{p^β} and $a^{p^\alpha}b$. Then by using the relations of the group G_2 , the resolutions for these pairs can be given as pictures \mathbf{K}_1 and \mathbf{K}_2 , respectively in Figure 5.

Now, let us also think the discs in the pictures \mathbf{K}_1 and \mathbf{K}_2 . To prove this theorem, we need to count the exponent sums of the discs in these pictures. So let us calculate the number of R_1 -discs, R_2 -discs and R_3 -discs in \mathbf{K}_1 , \mathbf{K}_2




Figure 5

where $R_1 : a^{p^\alpha} = 1$, $R_2 : b^{p^\beta} = 1$ and $R_3 : ab = ba^{1+p^{\alpha-\gamma}}$. Here, it is seen that

$$\exp_{R_1}(\mathbf{K}_1) = \frac{(1 + p^{\alpha-\gamma})^{p^\beta} - 1}{p^\alpha},$$

$$\exp_{R_1}(\mathbf{K}_2) = \frac{p^\alpha(1 + p^{\alpha-\gamma})}{p^\alpha} - 1 = p^{\alpha-\gamma},$$

$$\exp_{R_2}(\mathbf{K}_1) = 1 - 1 = 0,$$

$$\exp_{R_3}(\mathbf{K}_1) = 1 + (1 + p^{\alpha-\gamma}) + (1 + p^{\alpha-\gamma})^2 + \dots + (1 + p^{\alpha-\gamma})^{p^\beta-1} = \frac{(1 + p^{\alpha-\gamma})^{p^\beta} - 1}{p^{\alpha-\gamma}},$$

$$\exp_{R_3}(\mathbf{K}_2) = p^\alpha$$

and for the q -Cockcroft property to be held for some q , we need to have

$$\exp_{R_1}(\mathbf{K}_1) \equiv 0 \pmod{q} \Leftrightarrow \frac{(1 + p^{\alpha-\gamma})^{p^\beta} - 1}{p^\alpha} \equiv 0 \pmod{q},$$

$$\exp_{R_1}(\mathbf{K}_2) \equiv 0 \pmod{q} \Leftrightarrow p^{\alpha-\gamma} \equiv 0 \pmod{q},$$

$$\exp_{R_3}(\mathbf{K}_1) \equiv 0 \pmod{q} \Leftrightarrow \frac{(1 + p^{\alpha-\gamma})^{p^\beta} - 1}{p^{\alpha-\gamma}} \equiv 0 \pmod{q},$$

$$\exp_{R_3}(\mathbf{K}_2) \equiv 0 \pmod{p} \Leftrightarrow p^\alpha \equiv 0 \pmod{q}.$$

Here let us denote $\frac{(1+p^{\alpha-\gamma})^{p^\beta}-1}{p^\alpha}$ by A and $\frac{(1+p^{\alpha-\gamma})^{p^\beta}-1}{p^{\alpha-\gamma}}$ by B .

Therefore, since we have

$$\begin{aligned} (1 + p^{\alpha-\gamma})^{p^\beta} - 1 &= p^\beta p^{\alpha-\gamma} + \frac{1}{2} p^\beta (p^\beta - 1) p^{2(\alpha-\gamma)} + \frac{1}{6} p^\beta (p^\beta - 1) (p^\beta - 2) p^{3(\alpha-\gamma)} \\ &+ \dots + p^{p^\beta(\alpha-\gamma)} \end{aligned}$$

then we get that

$$\begin{aligned} A = p^{\beta-\gamma} &+ \frac{1}{2}p^{\beta}(p^{\beta}-1)p^{(\alpha-2\gamma)} + \frac{1}{6}p^{\beta}(p^{\beta}-1)(p^{\beta}-2)p^{(2\alpha-3\gamma)} \\ &+ \dots + p^{p^{\beta}(\alpha-\gamma)-\alpha} \end{aligned}$$

and

$$\begin{aligned} B = p^{\beta} &+ \frac{1}{2}p^{\beta}(p^{\beta}-1)p^{(\alpha-\gamma)} + \frac{1}{6}p^{\beta}(p^{\beta}-1)(p^{\beta}-2)p^{(2\alpha-2\gamma)} \\ &+ \dots + p^{p^{\beta}(\alpha-\gamma)-\alpha+\gamma}. \end{aligned}$$

Finally, we can see that $\pi_2(\mathcal{P}_{G_2})$ consists of the pictures \mathbf{K}_1 , \mathbf{K}_2 , \mathbf{C}_1 and \mathbf{C}_3 . Thus in order to get q -Cockcroft property, we must calculate the exponent sums of the discs in these pictures. Then, by using the above arguments, in order to get q -Cockcroft property for some prime q , we must have

$$\begin{aligned} A &\equiv 0 \pmod{q}, \quad p^{\alpha-\gamma} \equiv 0 \pmod{q}, \\ B &\equiv 0 \pmod{q}, \quad p^{\alpha} \equiv 0 \pmod{q}. \end{aligned}$$

Then by Theorem 1.1 we can say that the group G_2 is efficient if and only if it is q -Cockcroft for some prime q . Here since we have $\alpha \geq 2\gamma$ and $\beta > \gamma \geq 1$, then we choose $p = q$. So we get that the group G_2 presented by (5) is q -Cockcroft. This says that G_2 is efficient.

Remark 3.3. *We realised that we take $\beta > \gamma \geq 1$. If we take $\beta \geq \gamma \geq 1$, then we may have $\beta = \gamma$. This says that A is not equivalent to 0 by the modulo q for some prime q .*

Remark 3.4. *By using similar argumets as in Remark 3.2, since the group G_2 presented by (5) has non-negative deficiency and it is efficient as a group, then it is also efficient as a semigroup. So we deduce that the semigroup related to the certain group presentation (3) is also efficient.*

References

- [1] Arjomandfar, A., Campbell, C. M., Doostie, H., *Semigroups related to certain group presentations*, Semigroup Forum, Volume 85, Issue 3, (2012), 533-539.
- [2] Ates, F., Cevik, A. S., *The p -Cockcroft Property of Central Extensions of Groups II*, Monatshefte fr Math., 150 (2007), 181-191.

- [3] Ayık, H., Kuyucu, F., Vatansever, B., *On Semigroup Presentations and Efficiency*, Semigroup Forum, Vol. 65, (2002) 329335.
- [4] Baik, Y.G., Pride, S.J. *On the Efficiency of Coxeter Groups*, Bull. of the London Math. Soc. **29** (1997), 32-36.
- [5] Baik, Y.G., Harlander, j., Pride, S.J., *The Geometry of Group Extensions*, Journal of Group Theory **1**(4) (1998), 396-416.
- [6] Bacon, M.R., Kappe, L.C., *The nonabelian tensor square of a 2-generator p -group of class 2*. Arch. Math. (Basel) **61**, (1993) 508516.
- [7] Bogley W., A., and S., J., Pride, *Calculating generators of π_2* , in *Two-Dimensional Homotopy and Combinatorial Group Theory*, edited by C. Hog-Angeloni, W. Metzler, A. Sieradski, C.U. Press, 1993, pp. 157–188.
- [8] Campbell, C.M., Mitchell, J.D., Ruskuc, N., *On defining groups efficiently without using inverses*, Math. Proc. Cambridge Philos. Soc., **133** (2002), 31-36.
- [9] Çevik, A.S., *The p -Cockcroft Property of Central Extensions of Groups*, Comm. Algebra **29**(3) (2001), 1085-1094.
- [10] Çevik, A.S., *Minimal but inefficient presentations of the semidirect products of some monoids*, Semigroup Forum **66**, 1–17 (2003).
- [11] Dyer, M.N., *Cockcroft 2-Complexes*, preprint, University of Oregon, 1992.
- [12] Epstein, D.B.A., *Finite presentations of groups and 3-manifolds*, Quart. J. Math. **12**(2), 205–212 (1961).
- [13] Gilbert, N.D., Howie, J., *Threshold Subgroups for Cockcroft 2-Complexes*, Communications in Algebra **23**(1) (1995), 255-275.
- [14] Gilbert, N.D., Howie, J., *Cockcroft Properties of Graphs of 2-Complexes*, Proc. Royal Soc. of Edinburgh Section A-Mathematics **124**(2) (1994), 363-369.
- [15] Harlander, J., *Minimal Cockcroft Subgroups*, Glasgow Journal of Math. **36** (1994), 87-90.
- [16] Kappe, L.C., Sarmin, N., Visscher, M., *Two generator two-groups of class two and their non-Abelian tensor squares*. Glasg. Math. J. **41**, (1999), 417430.

- [17] Kilgour, C.W., Pride, S.J., *Cockcroft presentations*, *J. Pure Appl. Alg.* **106**(3), 275–295 (1996).
- [18] Pride, S.J., *Identities among relations of group presentations*, in *Group Theory from a Geometrical Viewpoint, Trieste 1990*, edited by E. Ghys, A. Haefliger, A. Verjovsky, editors, World Sci. Pub., 1991, pp. 687–717.

Firat Ateş,
Department of Mathematics,
Balikesir University,
10145 Balikesir, Turkiye
Email: frat@balikesir.edu.tr