

The Flora and Vegetation of Kunduz Forests (Vezirköprü / Samsun)

Fazıl ÖZEN

Balıkesir University, Faculty of Arts and Sciences, Department of Biology, 10100 Balıkesir - TURKEY

Mahmut KILINÇ

Ondokuz Mayıs University, Faculty of Arts and Sciences, Department of Biology, 55139 Samsun - TURKEY

Received: 29.11.2000

Accepted: 05.04.2002

Abstract: The research area, Kunduz Forests, is located on Kunduz Mountain, which is about 35 km south-west of Vezirköprü (Samsun). This area is in the A5 square in the grid scheme of Davis and is in the Euxine sector of the Euro-Siberian floristic region phytogeographically. The distribution of 306 specific and infraspecific taxa forming the flora of the research area according to floristic regions is as follows: Euro-Siberian 28.43%, Mediterranean 11.44% and Irano-Turanian 4.90%. The percentage of taxa found in more than one floristic region is 55.23%. Twenty-five of the 306 taxa found in the study area are given as new floristic records for the A5 square in the Flora of Turkey.

Besides the dominant forest vegetation type, there is also degraded forest vegetation in the research area. These vegetation types are represented by *Carpino-Quercetum cerridis* Kutbay & Kilinç 1995, *Daphno-Pinetum sylvestris* Kutbay & Kilinç 1995, *Junipero-Pinetum nigrae* Özgen & Kilinç ass. nov. and *Galio-Fagetum orientalis* Özgen & Kilinç ass. nov. associations. The last two out of the four associations mentioned here are determined to be new associations.

Key Words: Kunduz Mountain, Vezirköprü, Samsun, Flora, Vegetation.

Kunduz Ormanlarının (Vezirköprü / Samsun) Flora ve Vejetasyonu

Özet: Araştırma alanı olarak seçilen Kunduz Ormanları, Vezirköprü'nün (Samsun) 35 kilometre kadar güneybatısındaki Kunduz Dağı'nda bulunmaktadır. Davis'in kareleme sistemine göre A5 karesinde bulunan araştırma alanı, bitki coğrafyası bakımından Avrupa-Sibirya floristik bölgesinin Öksin Sektöründe yer almaktadır. Flora oluşturulan 306 tür ve türaltı seviyesindeki taksonun floristik bölgelere göre dağılımı şöyledir: Avrupa-Sibirya % 28,43, Akdeniz % 11,44, İran-Turan % 4,90 ve birden fazla bölgeler % 55,23. Bu çalışmada yer alan taksonlardan 25 adedi A5 karesi için yeni kayıt olarak verilmektedir.

Araştırma alanında hakim vejetasyon tipi olan orman vejetasyonunun yanısıra, bozuk orman vejetasyon tipi de bulunmaktadır. Bu vejetasyon tipleri *Carpino-Quercetum cerridis* Kutbay & Kilinç 1995, *Daphno-Pinetum sylvestris* Kutbay & Kilinç 1995, *Junipero-Pinetum nigrae* Özgen & Kilinç ass. nov. ve *Galio-Fagetum orientalis* Özgen & Kilinç Özgen & Kilinç ass. nov. bitki birlükleri ile temsil edilmektedir. Bu dört bitki birlüğinden son ikisi bilim dünyası için yenidir.

Anahtar Sözcükler: Kunduz Dağı, Vezirköprü, Samsun, Flora, Vejetasyon.

Introduction

In this study, the floristical and phytosociological aspects of Kunduz Forests, situated on Kunduz Mountain, and the surrounding area were studied. Although this area is one of the well-protected regions in Turkey, there has, except for those carried out in nearby regions (Özen & Kilinç, 1995 a, b and c; Kilinç, 1985; Kutbay & Kilinç, 1995 a and b; Karaer & Kilinç, 1993; Kilinç & Özgen, 1988), been no study on the vegetation and flora of the area. This area is of interest because there are some

Mediterranean floristic elements in the region. The aims of this study were to provide insight into the floristic and phytosociological structure of this area and to compare the data gained in the present study with those already known for some neighbouring localities.

Description of the Research Area

Kunduz Forests are located in the Central Black Sea area of the Black Sea region and are about 35 km south-

west of Vezirköprü (Samsun) (Figure 1). The altitude of the area varies between 400 and 1783 m. It is situated in the Euro-Siberian floristic region (Akman, 1993) and in the A5 square in Davis' grid scheme (Davis, 1965-1985; Davis et al., 1988). The geological structure of the region consists of Mesozoic, Upper Cretaceous and Metamorphic series (undifferentiated) (Erentöz, 1962). There are three large soil groups in the study area, namely grey brown podzolic soils, brown forest soils and chestnut soils (Anonymous, 1984). "Semiarid Mediterranean" bioclimate and the "Cold Winter Mediterranean" subsection of this bioclimate are seen in Vezirköprü according to the method of Emberger (Emberger, 1952). The precipitation regime in the study area is East Mediterranean Type 2 (Sp.W.A.S) (Akman, 1993).

Materials and Methods

The research area was visited during 1993-1996. Plants were identified using the Flora of Turkey (Davis, 1965-1985 and Davis et al., 1988). Plant specimens were kept at the herbarium of Balıkesir University (BAUH). The following strategy was used in order to list the flora: station, habitat characteristics, direction, altitude, collecting date, collector, specimen number,

floristic region of species and whether it is endemic or not. To avoid repetition "A5 Samsun: Vezirköprü" is not used each time. The plant list is given according to the rank of the Flora of Turkey. To indicate culture taxa and newly recorded taxa for the A5 square in Turkey, one asterisk and two asterisks are used, respectively. To identify new records for the A5 square in Turkey previous studies were used (Özen & Kılıç, 1995; Kutbay & Kılıç, 1995 a; Karaer & Kılıç, 1993; Kılıç & Özen, 1988; Kılıç & Özen, 1993; Özen, 2000; Donner, 1985, 1987 and 2000; Yıldırımlı, 1997 and Yıldırımlı, 1999).

The vegetation studies were carried out according to the Braun-Blanquet (1932) method. The plant associations were named and classified according to Syntaxonomic Nomenclatural Rules (Weber et al., 2000). During the classification of the vegetation, previous studies were used (Özen & Kılıç, 1995; Kılıç, 1985; Kutbay & Kılıç, 1995; Quezel et al., 1980; Akman et al., 1978, 1979, 1983 a and b and 1987; Akman, 1972 and 1973; Akman & Ketenoglu, 1978; Akman & Aydoğdu, 1986; Ocakverdi, 1987; Gemici, 1988; Tatlı, 1987; Ekim & Akman, 1990; Düzenli, 1989; Akman & İlarslan, 1983; Yurdakulol, 1981). Using Sorenson's similarity formula (Sorenson, 1948), the associations in the present study were compared with those given in the Black Sea region.




Figure 1. Map of study area

Results

Flora

This study was carried out on about 500 specimens. Two hundred and one species, 68 subspecies and 37

| | | |
|---|---|---|
| PTERIDOPHYTA | Pinus sylvestris L. | varieties, belonging to 190 genera and 70 families, were found in the research area. Among all the taxa examined, four were culture plants. |
| SPHENOPSIDA | Between Ovacık and deer production area, <i>Pinus sylvestris</i> forest, south-eastern slopes, 1150 m, 7.6.1995, F. Özen 1490, Euro-Siberian element. | BRASSICACEAE (Cruciferae) |
| EQUISETACEAE | | <i>Sinapis arvensis</i> L. |
| <i>Equisetum palustre</i> L. | Around deer production area, meadows, 1350 m, 7.6.1995, F. Özen 1596. | |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, streamside, 1200 m, 7.6.1995, F. Özen 1606. | <i>P. nigra</i> Arn. subsp. <i>pallasiana</i> (Lamb.) Holmboe | (**) <i>Crambe orientalis</i> L. var. <i>orientalis</i> |
| <i>E. arvense</i> L. | Around Soruk Village, north-eastern slopes, 1000 m, 8.6.1995, F. Özen 1499. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1250 m, 8.6.1995, F. Özen 1656, Irano-Turanian element. |
| HYPOLEPIDACEAE | CUPRESSACEAE | <i>Cardaria draba</i> (L.) Desv. subsp. <i>draba</i> |
| <i>Pteridium aquilinum</i> (L.) Kuhn | <i>Juniperus oxycedrus</i> L. subsp. <i>oxycedrus</i> | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1250 m, 8.6.1995, F. Özen 1657. |
| Around Karaköy Village, <i>Pinus sylvestris</i> forest clearings, 1000 m, 8.6.1995, F. Özen 1634. | Around Yukarı Narlı Village, <i>Pinus nigra</i> - <i>Juniperus oxycedrus</i> association, northern slopes, 400 m, 18.6.1993, F. Özen 1190. | <i>Thlaspi orbiculatum</i> Steven |
| ATHYRIACEAE | ANGIOSPERMAE | Around Soruk Village, <i>Pinus sylvestris</i> forest, 1400 m, 9.6.1995, F. Özen 1601. |
| <i>Cystopteris fragilis</i> (L.) Bernh. | DICOTYLEDONAE | |
| Around fire observation tower, <i>Fagus orientalis</i> - <i>Galium odoratum</i> association, northern slopes, 1720 m, 1.8.1994, F. Özen 1374. | RANUNCULACEAE | T. <i>perfoliatum</i> L. |
| ASPIDIACEAE | <i>Ranunculus repens</i> L. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1250 m, 8.6.1995, F. Özen 1670. |
| <i>Polystichum setiferum</i> (Forsk.) Woyn. | Between Ovacık and Soruk Villages, <i>Pinus sylvestris</i> forest clearings, 1100 m, 7.6.1995, F. Özen 1611. | T. <i>annuum</i> K.Koch |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, <i>Pinus sylvestris</i> forest, south-western slopes, 1150 m, 7.6.1995, Özen, 1484. | <i>R. illyricus</i> L. subsp. <i>illyricus</i> | Around deer production area, <i>Pinus sylvestris</i> forest, 1350 m, 7.6.1995, F. Özen 1580. |
| SPERMATOPHYTA | Around deer production area, <i>Pinus sylvestris</i> forest clearings, 1300 m, 7.6.1995, F. Özen 1570. | <i>Capsella bursa-pastoris</i> (L.) Medik. |
| GYMNOSPERMAE | R. <i>arvensis</i> L. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1250 m, 23.7.1994, F. Özen 1282. |
| PINACEAE | Around deer production area, meadows, 1300 m, 7.6.1995, F. Özen 1547. | <i>Erophila verna</i> (L.) Chevall. subsp. <i>verna</i> |
| <i>Abies nordmanniana</i> (Steven) Spach subsp. <i>bormmuelleriana</i> (Mittf.) Coode & Cullen | (**) R. <i>sphaerospermus</i> Boiss. & Blanche | Between Ovacık and deer production area, <i>Pinus sylvestris</i> forest clearings, 1300 m, 1.6.1995, F. Özen 1609. |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, northern slopes, 1250 m, 29.9.1993, F. Özen 1195, Endemic, Euxine element. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, ponds, 1250 m, 10.6.1995, F. Özen 1605. | <i>Turritis laxa</i> (Sibth. & Sm.) Hayek |
| PAPAVERACEAE | PAPAVERACEAE | Between Ovacık and deer production area, <i>Pinus sylvestris</i> forest clearings, 1300 m, 1.6.1995, F. Özen 1642. |
| <i>Papaver rhoeas</i> L. | <i>Papaver rhoeas</i> L. | <i>Cardamine bulbifera</i> (L.) Crantz |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1150 m, 8.6.1995, F. Özen 1683. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1150 m, 8.6.1995, F. Özen 1494, Euro-Siberian element. | Between Ovacık and Karaköy Villages, <i>Fagus orientalis</i> - <i>Galium odoratum</i> association, north-eastern slopes, 1150 m, 9.6.1995, F. Özen 1494, Euro-Siberian element. |

| | | |
|---|--|--|
| C. quingifolia (M.Bieb.) Schmalh. | POLYGALACEAE | POLYGONACEAE |
| Around fire observation tower, <i>Fagus orientalis</i> - <i>Galium odoratum</i> association, north-western slopes, 1700 m, 1.8.1996, F. Özen 1701, Euro-Siberian element. | <i>Polygala pruinosa</i> Boiss. subsp. <i>pruinosa</i> | (**) <i>Polygonum arenastrum</i> Bor. |
| <i>Sisymbrium loeselii</i> L. | Around Hamidiye Village, field margins, south-eastern slopes, 600 m, 7.6.1995, F. Özen 1564. | Around fire observation tower, rocks, 1700 m, 1.8.1995, F. Özen 1455. |
| Between Ovacık and deer production area, <i>Pinus sylvestris</i> forest clearings, 1300 m, 1.6.1995, F. Özen 1610. | <i>P. anatolica</i> Boiss. & Heldr. | <i>P. bellardii</i> All. |
| RESEDACEAE | Between Mezraa and Sofular Village, <i>Pinus nigra</i> - <i>Juniperus oxycedrus</i> association, southern slopes, 800 m, 1.6.1995, F. Özen 1525. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1250 m, 20.7.1995, F. Özen 1297. |
| Reseda lutea L. var. <i>nutans</i> Boiss. | CARYOPHYLLACEAE | Rumex acetosella L. |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, <i>Pinus sylvestris</i> forest clearings, 1200 m, 9.6.1995, F. Özen 1661. | (**) <i>Arenaria leptoclados</i> (Rchb.) Guss. | Around fire observation tower, <i>Fagus orientalis</i> forest, north-eastern slopes, 1700 m, 1.8.1994, F. Özen 1393. |
| R. luteola L. | Around fire observation tower, rocks, 1700 m, 1.8.1995, F. Özen 1457. | (**) <i>R. alpinus</i> L. |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1200 m, 20.7.1994, F. Özen 1314. | <i>A. serpyllifolia</i> L. | Around Tahtaköprü Village, streamside, 1000 m, 10.6.1995, F. Özen 1628. |
| CISTACEAE | Between Soruk Village-fire observation tower, <i>Fagus orientalis</i> forest, north-eastern slopes, 1500 m, 31.7.1994, F. Özen 1401. | R. patientia L. |
| <i>Cistus creticus</i> L. | <i>Stellaria holostea</i> L. | Around Sofular Village, <i>Fagus orientalis</i> forest clearings, 1150 m, 20.7.1994, F. Özen 1202. |
| Around Sofular Village, <i>Pinus nigra</i> forest, southern slopes, 1160 m, 31.7.1994, F. Özen 1418, Omni-Mediterranean element. | Around Göltürkmen Village, <i>Quercus petraea</i> forest, 950 m, 8.6.1995, F. Özen 1496, Euro-Siberian element. | CHENOPODIACEAE |
| <i>C. laurifolius</i> L. | <i>Moenchia mantica</i> (L.) Bartl. subsp. <i>mantica</i> | <i>Chenopodium botrys</i> L. |
| Around Sofular Village, <i>Pinus nigra</i> forest, southern slopes, 1130 m, 31.7.1994, F. Özen 1417, Mediterranean element. | Around deer production area, <i>Pinus sylvestris</i> forest clearings, 1350 m, 7.6.1995, F. Özen 1602. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1180 m, 20.7.1994, F. Özen 1275. |
| <i>Helianthemum nummularium</i> (L.) Mill. subsp. <i>nummularium</i> | <i>Dianthus armeria</i> L. subsp. <i>armeria</i> | <i>C. murale</i> L. |
| Around Göltürkmen Village, <i>Quercus cerris</i> - <i>Carpinus orientalis</i> association, 1040 m, 31.7.1994, F. Özen 1421. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1200 m, 22.7.1994, F. Özen 1277, Euro-Siberian element. | Around Soruk Village, <i>Pinus sylvestris</i> forest clearings, south-eastern slopes, 1400 m, 31.7.1994, F. Özen 1444. |
| <i>Fumana thymifolia</i> (L.) Verl. var. <i>viridis</i> (Ten.) Boiss. | <i>D. carmelitarum</i> Reut. ex Boiss. | <i>Beta corolliflora</i> Zosimovic ex Buttler |
| Around Gölalan Village, <i>Pinus nigra</i> - <i>Juniperus oxycedrus</i> association, 900 m, 8.6.1995, F. Özen 1506, Mediterranean element. | Around Gölalan Village, <i>Fagus orientalis</i> forest, 1110 m, 21.7.1994, F. Özen 1233, Endemic, Euxine element. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, <i>Pinus sylvestris</i> forest clearings, 1100 m, 8.6.1995, F. Özen 1658, Iran-Turanian element. |
| VIOLACEAE | <i>Silene vulgaris</i> (Moench) Garcke var. <i>vulgaris</i> | TAMARICACEAE |
| <i>Viola odorata</i> L. | Summit of Kunduz Mountain, <i>Fagus orientalis</i> - <i>Galium odoratum</i> association, 1720 m, 31.7.1994, F. Özen 1387. | <i>Myricaria germanica</i> (L.) Desv. |
| Around deer production area, <i>Pinus sylvestris</i> - <i>Daphne pontica</i> association, south-eastern slopes, 1350 m, 25.9.1993, F. Özen 1194. | <i>Agrostemma githago</i> L. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, streamside, 1200 m, 22.7.1994, F. Özen 1316. |
| ILLECEBRACEAE | Around Göltürkmen Village, <i>Fagus orientalis</i> forest, 1100 m, 20.7.1994, F. Özen 1212. | HYPERICACEAE (Guttiferae) |
| <i>Herniaria glabra</i> L. | <i>Herniaria glabra</i> L. | <i>Hypericum androsaemum</i> L. |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1200 m, 20.7.1994, F. Özen 1283. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1200 m, 20.7.1994, F. Özen 1250. | |

- H. linarioides** Bosse
Summit of Kunduz Mountain, *Fagus orientalis*-*Galium odoratum* association, 1740 m, 1.8.1995, F. Özen 1385.
- (**)H. atomarium Boiss.
Around Yukarı Narlı Village, field margins, 450 m, 9.6.1995, F. Özen 1562, East Mediterranean element.
- (**)H. bithynicum Boiss.
Around Sırbaşmak Village, roadsides, 400 m, 1.6.1995, F. Özen 1626, Euxine element.
- (**)H. olympicum L. subsp. *olympicum*
Around Gölköy-Asarcık forest store, *Pinus nigra* forest clearings, 1080 m, 13.7.1994, F. Özen 1469, East Mediterranean element.
- H. perforatum L.
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1300 m, 13.7.1994, F. Özen 1312.
- TILIACEAE**
Tilia rubra DC. subsp. *caucasica* (Rupr.) V. Engl.
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1200 m, 8.6.1995, Özen, 1660, Euxine element.
- LINACEAE**
Linum flavum L. subsp. *scabrinerve* (P.H.Davis) P.H.Davis
Around Hamidiye Village, *Pinus nigra* forest, south-eastern slopes, 750 m, 8.6.1995, F. Özen 1517, Endemic, Iran-Turanian element.
- L. corymbulosum Rchb.
Around Hamidiye Village, *Pinus nigra* forest, south-eastern slopes, 750 m, 8.6.1995, F. Özen 1518, Mediterranean element.
- L. aroanium Boiss. & Orph.
Around Yukarı Narlı Village, field margins, 500 m, 8.6.1995, F. Özen 1559.
- GERANIACEAE**
Geranium pusillum Burm.f.
Around fire observation tower, rocks, 1700 m, 1.8.1994, F. Özen 1454.
- G. aspedoleoides Burm.f. subsp. *aspedoleoides*
Around deer production area, *Pinus sylvestris*-*Daphne pontica* association, eastern slopes, 1400 m, 2.8.1996, F. Özen 1705, Euro-Siberian element.
- G. pyrenaicum Burm.f.
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, meadows, 1250 m, 8.6.1995, F. Özen 1685.
- Erodium cicutarium (L.) L'Hér. subsp. *cicutarium*
Around Gölalan Village, *Fagus orientalis* forest clearings, 1150 m, 10.7.1994, F. Özen 1223.
- SIMAROUBACEAE**
(*)*Ailanthus altissima* (Mill.) Swingle
Around Yukarı Narlı Village, roadsides, 500 m, 8.6.1995, F. Özen 1612.
- ACERACEAE**
Acer campestre L. subsp. *leiocarpum* (Opiz) Pax
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, *Fagus orientalis* forest, 1030 m, 27.7.1994, F. Özen 1364, Euro-Siberian element.
- RHAMNACEAE**
Paliurus spina-christi Mill.
Around Sırbaşmak Village, 400 m, 8.6.1995, F. Özen 1618.
- ANACARDIACEAE**
Rhus coraria L.
Around Bakla Village, *Pinus nigra*-*Juniperus oxycedrus* association, 800 m, 1.8.1995, F. Özen 1438.
- (*)*Pistacia vera* L.
Around Sırbaşmak Village, field margins, 400 m, 8.6.1995, F. Özen 1616, Iran-Turanian element.
- P. terebinthus* L. subsp. *terebinthus*
Around Sırbaşmak Village, field margins, 450 m, 8.6.1995, F. Özen 1615, Mediterranean element.
- FABACEAE** (Leguminosae)
Genista lydia Boiss. var. *lydia*
Around Göltürkmen Village, *Quercus cerris*-*Carpinus orientalis* association, north-eastern slopes, 1000 m, 8.6.1995, F. Özen 1495.
- Argyrolobium biebersteinii** Ball
Around Mezraa, *Pinus nigra*-*Juniperus oxycedrus* association, south-western slopes, 800 m, 9.6.1995, F. Özen 1511.
- Astragalus angustifolius* Lam. subsp. *angustifolius* var. *angustifolius*
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, *Pinus sylvestris* forest clearings, 1150 m, 8.6.1995, F. Özen 1655.
- Psoralea bituminosa** L.
Around Mezraa, *Pinus nigra*-*Juniperus oxycedrus* association, south-western slopes, 750 m, 7.6.1995, F. Özen 1514, Mediterranean element.
- Vicia crocea* (Desf.) B.Fedtsch.
Around Zanbal Village, *Fagus orientalis*-*Galium odoratum* association, north-eastern slopes, 1300 m, 20.7.1994, F. Özen 1328, Hyrcano-Euxine element.
- (**)V. villosa Roth subsp. *eriocarpa* (Hausskn.) P.W.Ball
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1250 m, 8.6.1995, F. Özen 1681.
- (**)V. villosa Roth subsp. *dasyocarpa* (Ten.) Cav.
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1200 m, 22.7.1994, F. Özen 1278.
- V. peregrina L.
Around Sofular Village, field margins, 1100 m, 21.7.1994, F. Özen 1213.
- V. lutea L. var. *hirta* (Balb.) Lois.
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1250 m, 15.6.1995, F. Özen 1672.
- V. sativa L. subsp. *nigra* (L.) Ehrh. var. *nigra*
Around Yukarı Narlı Village, field margins, 500 m, 8.6.1995, F. Özen 1556.
- V. sativa L. subsp. *nigra* (L.) Ehrh. var. *segetalis* (Thuill.) Ser. ex DC.
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1200 m, 22.7.1994, F. Özen 1296.

| | | |
|--|--|--|
| <i>Lathyrus tukhtensis</i> Czecott | (**)T. <i>purpureum</i> Loisel var. <i>purpureum</i> | ROSACEAE |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, <i>Pinus sylvestris</i> forest clearings, north-western slopes, 1250 m, 8.6.1995, F. Özen 1680, Endemic. | Around deer production area, <i>Pinus sylvestris</i> forest, north-eastern slopes, 1400 m, 7.6.1995, F. Özen 1599. | Filipendula vulgaris Moench |
| <i>L. pratensis</i> L. | <i>Melilotus officinalis</i> (L.) Desr. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1250 m, 21.6.1995, F. Özen 1676, Euro-Siberian element. |
| Around Karaköy Village, <i>Fagus orientalis</i> forest, south-western slopes, 1150 m, 8.6.1995, F. Özen 1629. | Around Gölalan Village, field margins, 1110 m, 22.7.1994, F. Özen 1206. | <i>Rubus discolor</i> Weihe & Nees |
| <i>L. laxiflorus</i> (Desf.) Kuntze subsp. <i>laxiflorus</i> | <i>Medicago lupulina</i> L. | Around Gölalan Village, <i>Fagus orientalis</i> forest, 1100 m, 22.7.1994, F. Özen 1231. |
| Around Gölalan Village, <i>Fagus orientalis</i> - <i>Galium odoratum</i> association, 1720 m, 31.7.1994, F. Özen 1380. | Around Soruk Village, <i>Pinus sylvestris</i> forest clearings, 1400 m, 31.7.1994, F. Özen 1446. | <i>R. canescens</i> DC. var. <i>canescens</i> |
| (**)L. <i>laxiflorus</i> (Desf.) Kuntze subsp. <i>angustifolius</i> | <i>Dorycnium graecum</i> (L.) Ser. | Around Zanbal Village, 1150 m, 22.7.1994, F. Özen 1284, Euro-Siberian element. |
| Around deer production area, <i>Pinus sylvestris</i> - <i>Daphne pontica</i> association, eastern slopes, 1350 m, 2.7.1994, F. Özen 1355, Endemic, East Mediterranean element. | Around Karaköy Village, <i>Fagus orientalis</i> forest, western slopes, 1150 m, 10.7.1995, F. Özen 1254, Euxine element. | <i>R. hirtus</i> Waldst. & Kit. |
| <i>L. aphaca</i> L. var. <i>pseudoaphaca</i> (Boiss.) P.H.Davis | D. <i>pentaphyllum</i> Scop. subsp. <i>anatolicum</i> (Boiss.) Gams | Between Ovacık and deer production area, <i>Fagus orientalis</i> forest, north-eastern slopes, 1270 m, 9.6.1995, F. Özen 1488, Euro-Siberian element. |
| Around Mezraa, <i>Pinus nigra-Juniperus oxycedrus</i> association, south-eastern slopes, 750 m, 8.6.1995, F. Özen 1527, East Mediterranean element. | Around Mezraa, <i>Pinus nigra-Juniperus oxycedrus</i> association, south-eastern slopes, 750 m, 8.6.1995, F. Özen 1521. | <i>Potentilla argentea</i> L. |
| <i>Ononis pusilla</i> L. | D. <i>pentaphyllum</i> Scop. subsp. <i>herbaceum</i> (Vill.) Rouy | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, <i>Pinus sylvestris</i> forest, 1300 m, 20.6.1995, F. Özen 1666. |
| Around Yukarı Narlı Village, field margins, 500 m, 8.6.1995, F. Özen 1558, Mediterranean element. | Around Yukarı Narlı Village, field margins, 500 m, 9.6.1995, F. Özen 1567. | <i>P. inclinata</i> Vill. |
| <i>Trifolium campestre</i> Schreb. | <i>Lotus angustissimus</i> L. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, <i>Pinus sylvestris</i> forest, 1300 m, 20.6.1995, F. Özen 1665. |
| Around Soruk Village, <i>Pinus sylvestris</i> forest clearings, north-eastern slopes, 1400 m, F. Özen 1445. | Around deer production area, <i>Pinus sylvestris</i> - <i>Daphne pontica</i> association, north-eastern slopes, 1375 m, 20.7.1994, F. Özen 1334. | <i>P. recta</i> L. Grup B |
| <i>T. medium</i> L. var. <i>medium</i> | <i>L. corniculatus</i> L. var. <i>corniculatus</i> | Around deer production area, <i>Pinus sylvestris</i> forest, 1250 m, 7.6.1995, F. Özen 1573. |
| Around Sofular Village, <i>Pinus nigra</i> forest, southern slopes, 1130 m, 1.8.1994, F. Özen 1415. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1250 m, 8.6.1995, F. Özen 1667. | <i>P. reptans</i> L. |
| <i>T. caudatum</i> Boiss. | <i>L. corniculatus</i> L. var. <i>tenuifolius</i> L. | Ovacık Village, around Burhan Zorlu Education Centre belonging Ministry of Forestry, field margins, 1300 m, 23.7.1994, F. Özen 1311. |
| Around fire observation tower, <i>Fagus orientalis</i> forest, north-eastern slopes, 1700 m, 31.7.1994, F. Özen 1394, Endemic. | <i>L. corniculatus</i> L. var. <i>alpinus</i> Ser. | <i>Fragaria vesca</i> L. |
| <i>T. hirtum</i> All. | Around fire observation tower, rocks, 1700 m, 31.7.1994, F. Özen 1453. | Around Karaköy Village, <i>Fagus orientalis</i> forest, northern slopes, 1100 m, 13.7.1995, F. Özen 1265. |
| Around Sofular Village, <i>Fagus orientalis</i> forest clearings, eastern slopes, 1000 m, 13.7.1994, F. Özen 1218, Mediterranean element. | <i>Coronilla scorpioides</i> (L.) K.Koch | <i>Geum urbanum</i> L. |
| <i>C. varia</i> L. subsp. <i>varia</i> | Around Mezraa, <i>Pinus nigra-Juniperus oxycedrus</i> association, south-eastern slopes, 750 m, 7.6.1995, F. Özen 1513. | Around Karaköy Village, <i>Fagus orientalis</i> - <i>Galium odoratum</i> association, 1700 m, 31.7.1994, F. Özen 1382, Euro-Siberian element. |
| Around Gölalan Village, field margins, 1110 m, 22.7.1994, F. Özen 1205. | | |

Sanguisorba minor Scop. subsp.
muricata (Spach) Briq.

Around Yukarı Narlı Village, field margins, 550 m, 8.6.1995, F. Özen 1557.

(**) *Alchemilla surculosa* E.Fröhner

Around fire observation tower, *Fagus orientalis* forest, north-eastern slopes, 1675 m, 31.7.1994, F. Özen 1390, Hyrcano-Euxine (mountain) element.

(**) *A. lithophila* Juz.

Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1150 m, 20.6.1995, F. Özen 1653.

A. heterophylla Rothm.

Around deer production area, 1350 m, 18.6.1995, F. Özen 1603.

Rosa canina L.

Around deer production area, *Pinus sylvestris-Daphne pontica* association, eastern slopes, 1350 m, 19.7.1994, F. Özen 1349.

Pyracantha coccinea Roem.

Around deer production area, *Pinus sylvestris* forest clearings, 1350 m, 7.6.1995, F. Özen 1542.

Crataegus tanacetifolia (Lam.) Pers.

Around deer production area, *Pinus sylvestris-Daphne pontica* association, western slopes, 1330 m, 7.6.1995, F. Özen 1541, Endemic.

C. monogyna Jacq. subsp. *monogyna*

Around deer production area, *Pinus sylvestris* forest, 1250 m, 7.6.1995, F. Özen 1535.

C. microphylla K.Koch

Around Gölalan Village, *Fagus orientalis* forest, 1030 m, 20.7.1994, F. Özen 1365, Hyrcano-Euxine element.

Sorbus torminalis (L.) Crantz var. *pinnatifida* Boiss.

Around Göltürkmen Village, *Quercus cerris-Carpinus orientalis* association, north-eastern slopes, 1000 m, 8.6.1995, F. Özen 1503.

(*) *Pyrus communis* L. subsp. *sativa* (DC.) Hegi

Around Sirbaşmak Village, field margin, 400 m, 15.6.1995, F. Özen 1617.

LYTHRACEAE

Lythrum salicaria L.

Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, streamsides, 1150 m, 20.7.1994, F. Özen 1280, Euro-Siberian element.

ONAGRACEAE

Ciraea lutetiana L.

Around Göltürkmen Village, *Fagus orientalis* forest clearings, north-western slopes, 1100 m, 10.7.1994, F. Özen 1253.

Epilobium angustifolium L.

Between Ovacık and deer production area, *Pinus sylvestris* forest clearings, north-western slopes, 1400 m, 31.7.1994, F. Özen 1464.

E. parviflorum Schreb.

Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1180 m, 22.7.1994, F. Özen 1270.

E. hirsutum L.

Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1180 m, 22.7.1994, F. Özen 1272.

E. montanum L.

Around Göltürkmen Village, *Pinus sylvestris* forest clearings, 1150 m, 10.7.1995, F. Özen 1244, Euro-Siberian element.

E. lanceolatum Seb. & Mauri

Between Ovacık and deer production area, north-eastern slopes, 1250 m, 7.6.1995, F. Özen 1481.

E. palustre L.

Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1100 m, 22.7.1994, F. Özen 1271.

(**) *E. ponticum* Hausskn.

Around Karaköy Village, *Fagus orientalis* forest, north-western slopes, 1100 m, 11.7.1994, F. Özen 1256.

CRASSULACEAE

Sedum pallidum M.Bieb. var. *pallidum*

Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, *Pinus sylvestris* forest clearings, 1180 m, 22.7.1994, F. Özen 1285, Euxine element.

S. pallidum M.Bieb. var. *bithynicum* (Boiss.) Chamb.

Around fire observation tower, *Fagus orientalis-Galium odoratum* association, 1740 m, 31.7.1994, F. Özen 1369.

SAXIFRAGACEAE

Saxifraga cymbalaria L. var. *cymbalaria*

Around Karaköy Village, *Fagus orientalis* forest clearings, 1100 m, 13.7.1995, F. Özen 1262.

APIACEAE (Umbelliferae)

Sanicula europaea L.

Around deer production area, *Pinus sylvestris-Daphne pontica* association, northern slopes, 1350 m, 20.7.1994, F. Özen 1335, Euro-Siberian element.

Chaerophyllum temulum L.

Around Gölköy Village, *Fagus orientalis* forest clearings, 1150 m, 2.7.1994, F. Özen 1200, Euro-Siberian element.

Bifora radians M.Bieb.

Around Gölalan Village, field margins, 450 m, 8.6.1995, F. Özen 1625.

Seseli peucedanoides (M.Bieb.) Kosov Pol.

Around Hamidiye Village, *Pinus nigra-Juniperus oxycedrus* association, south-western slopes, 750 m, 10.6.1995, F. Özen 1531, Euro-Siberian element.

Trinia scabra Boiss. & Noe

Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, *Pinus sylvestris* forest clearings, 1250 m, 9.6.1995, F. Özen 1692, Endemic?, Irano-Turanian element.

(**) *Peucedanum chryseum* (Boiss. & Heldr.) Chamb.

Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1200 m, 22.7.1994, F. Özen 1322, Endemic.

Orlaya daucoides (L.) Greuter

Around deer production area, *Pinus sylvestris* forest, 1350 m, 7.6.1995, F. Özen 1550, Mediterranean element ?

Artemia squamata L.

Around Gölköy-Asarcık forest store, roadsides, 1080 m, 31.7.1994, F. Özen 1470.

| | | |
|---|---|---|
| ARALIACEAE | DIPSACACEAE | Achillea millefolium L. subsp. millefolium |
| <i>Hedera helix L.</i> | <i>Dipsacus laciniatus L.</i> | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1300 m, 23.7.1994, F. Özen 1647. |
| Around deer production area, <i>Pinus sylvestris</i> forest, 1250 m, 8.6.1995, F. Özen 1647. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1300 m, 23.7.1994, F. Özen 1318. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, <i>Pinus sylvestris</i> forest clearings, 1250 m, 8.6.1995, F. Özen 1679, Euro-Siberian element. |
| CORNACEAE | Scabiosa columbaria L. subsp. ochroleuca (L.) Celak. var. ochroleuca (L.) Coult. | A. coarctata Poir. |
| <i>Cornus sanguinea L. subsp. australis (C.A.Mey.) Jav.</i> | Around Gölköy, Village, roadsides, 1100 m, 6.7.1994, F. Özen 1229. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1180 m, 22.7.1994, F. Özen 1300. |
| Around Öğürlü Village, <i>Quercus cerris-Carpinus orientalis</i> association, south-western slopes, 1000 m, 29.7.1994, F. Özen 1426, Euro-Siberian element. | S. argentea L. | Tanacetum parthenium (L.) Sch.Bip. |
| <i>C. mas L.</i> | Around Karaköy Village, <i>Pinus nigra-Juniperus oxycedrus</i> association, 800 m, 31.7.1994, F. Özen 1439. | Around Gölköy Village, <i>Fagus orientalis</i> forest clearings, 1100 m, 20.7.1994, F. Özen 1225. |
| Around Sirbaşmak Village, 350 m, 8.6.1995, F. Özen, 1620, Euro-Siberian element? | S. micrantha Desf. | Arctium minus (Hill) Bernh. subsp. pubens (Bab.) Arénes |
| CAPRIFOLIACEAE | Around Gölköy Village, field margins, 500 m, 8.6.1995, F. Özen 1624. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, <i>Pinus sylvestris</i> forest clearings, 1200 m, 23.7.1994, F. Özen 1317, Euro-Siberian element. |
| <i>Sambucus ebulus L.</i> | ASTERACEAE (Compositae) | Cirsium pseudopersonata Boiss. & Bal. subsp. pseudopersonata |
| Around Gölalan Village, <i>Fagus orientalis</i> forest clearings, 1110 m, 3.7.1994, F. Özen 1227, Euro-Siberian element. | <i>Filago eriocephala Guss.</i> | Around Karaköy Village, <i>Pinus sylvestris</i> forest, 1150 m, 13.7.1994, F. Özen 1264, Endemic, Euxine element. |
| <i>S. nigra L.</i> | Around Mezraa, <i>Pinus nigra-Juniperus oxycedrus</i> association, south-western slopes, 800 m, 9.6.1995, F. Özen 1528, East Mediterranean element. | Centaurea hypoleuca DC. |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, <i>Pinus sylvestris</i> forest clearings, north-eastern slopes, 1130 m, 29.7.1994, F. Özen 1695, Euro-Siberian element. | <i>F. pyramidata L.</i> | Around Mezraa, <i>Pinus nigra-Juniperus oxycedrus</i> association, south-western slopes, 750 m, 7.6.1995, F. Özen 1515, Euxine element. |
| <i>Viburnum lantana L.</i> | Around fire observation tower, rocks, 1700 m, 1.8.1995, F. Özen 1458. | Cichorium intybus L. |
| Around Sofular Village, <i>Fagus orientalis</i> forest, 1030 m, 13.7.1994, F. Özen, 1367, Euro-Siberian element. | <i>Bellis perennis L.</i> | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1200 m, 22.7.1994, F. Özen 1291. |
| <i>Lonicera etrusca Santi var. etrusca</i> | Around deer production area, <i>Pinus sylvestris</i> forest, 1350 m, 7.6.1995, F. Özen 1588, Euro-Siberian element. | Tragopagon longirostris Bisch. ex Schultz var. longirostris |
| Around Sirbaşmak Village, field margins, 400 m, 8.6.1995, F. Özen 1619, Mediterranean element. | <i>Senecio vernalis Waldst. & Kit.</i> | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1250 m, 15.6.1995, F. Özen 1659. |
| VALERIANACEAE | Around fire observation tower, <i>Pinus sylvestris</i> forest clearings, south-western slopes, 1400 m, 31.7.1994, F. Özen 1449. | T. aureus Boiss. |
| <i>Valerianella locusta (L.) Laterr.</i> | <i>Tussilago farfara L.</i> | Around Mezraa, <i>Pinus nigra-Juniperus oxycedrus</i> association, south-eastern slopes, 700 m, 7.6.1995, F. Özen 1509, Endemic. |
| Around deer production area, <i>Pinus sylvestris</i> forest, north-eastern slopes, 1400 m, 9.6.1994, F. Özen 1577, Euro-Siberian element? | Around Karaköy Village, <i>Fagus orientalis</i> forest clearings, 1000 m, 8.6.1995, F. Özen 1633, Euro-Siberian element. | |
| MORINACEAE | <i>Anthemis tinctoria L. var. pallida DC.</i> | |
| <i>Morina persica L.</i> | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, <i>Pinus sylvestris</i> forest clearings, 1250 m, 8.6.1995, F. Özen 1675. | |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1100 m, 8.6.1996, F. Özen 1652, Irano-Turanian element. | | |

Pilosella hoppeana (H.Schult.) C.H. & F.W.Schultz subsp. *pilisquama* (NP.) Sell & West

Around fire observation tower, rocks, 1700 m, 1.8.1995, F. Özen 1452.

P. hoppeana (Schultes) C.H. & F.W.Schultz subsp. *testimonialis* (Naegli ex Peter) Sell & West

Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, *Pinus sylvestris* forest clearings, 1200 m, 22.7.1994, F. Özen 1290, Euro-Siberian element.

P. x auricloides (Lang) Sell & West

Around deer production area, *Pinus sylvestris* forest, 1300 m, 7.6.1995, F. Özen 1475.

Mycelis muralis (L.) Dum.

Around deer production area, *Pinus sylvestris*-*Daphne pontica* association, north-eastern slopes, 1350 m, 20.7.1994, F. Özen 1345, Euro-Siberian element.

CAMPANULACEAE

Campanula rapunculoides L. subsp. *rapunculoides*

Around Tahtaköprü Village, *Pinus nigra* forest clearings, southern slopes, 1125 m, 31.7.1994, F. Özen 1409, Euro-Siberian element.

C. glomerata L. subsp. *hispanica* (Witasek) Hayek

Around Soruk Village, *Fagus orientalis*-*Galium odoratum* association, 1720 m, 31.7.1994, F. Özen 1386, Euro-Siberian element.

C. saxonorum Gand.

Around fire observation tower, 1675 m, 31.7.1994, F. Özen 1451, Endemic, Iran-Turanian element?

Asyneuma rigidum (Willd.) Grossh. subsp. *rigidum*

Around Mezraa, *Pinus nigra*-*Juniperus oxycedrus* association, south-eastern slopes, 1000 m, 8.6.1995, F. Özen 1520, Iran-Turanian element.

Legousia speculum-veneris (L.) Chaix

Around Soruk Village, *Fagus orientalis*-*Galium odoratum* association, 1750 m, 31.7.1994, F. Özen 1381, Mediterranean element.

ERICACEAE

Moneses uniflora (L.) A.Gray

Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, *Fagus orientalis* forest, north-western slopes, 1300 m, 1.6.1995, F. Özen 1487.

Monotropa hypopithys L.

Around fire observation tower, *Fagus orientalis*-*Galium odoratum* association, north-western slopes, 1740 m, 29.7.1994, F. Özen 1368.

PRIMULACEAE

Primula vulgaris Huds. subsp. *vulgaris*

Around Soruk Village, *Fagus orientalis* forest, northern slopes, 1300 m, 3.8.1996, F. Özen 1707, Euro-Siberian element.

Cyclamen coum Mill. var. *coum*

Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, *Fagus orientalis* forest, north, 4.8.1996, F. Özen 1709.

Lysimachia verticillaris Spreng.

Around Karaköy Village, streamside, north-western slopes, 1100 m, 11.7.1995, F. Özen 1257, Hyrcano-Euxine element.

OLEACEAE

Jasminum fruticans L.

Around Yukarı Narlı Village, *Pinus nigra*-*Juniperus oxycedrus* association, 400 m, 4.8.1996, F. Özen 1710, Mediterranean element.

Fontanesia philliraeoides Labill. subsp. *philliraeoides*

Around Sırbaşmak Village, 450 m, 8.6.1995, F. Özen 1627, East Mediterranean element.

Fraxinus excelsior L. subsp. *excelsior*

Around fire observation tower, *Fagus orientalis*-*Galium odoratum* association, northern slopes, 1720 m, 30.7.1996, F. Özen 1700, Euro-Siberian element.

Ligustrum vulgare L.

Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1200 m, 8.6.1995, F. Özen 1691, Euro-Siberian element.

Phillyrea latifolia

Around Yukarı Narlı Village, 870 m, 31.7.1994, F. Özen 1434, Mediterranean element.

GENTIANACEAE

Centaurium erythraea Rafn subsp. *erythraea*

Around Karaköy Village, north-western slopes, 1150 m, 20.7.1994, F. Özen 1259, Euro-Siberian element.

Gentiana cruciata L.

Around Karaköy Village, north-western slopes, 1150 m, 20.7.1994, F. Özen 1267, Euro-Siberian element.

CONVOLVULACEAE

Convolvulus cantabrica L.

Around Mezraa, *Pinus nigra* forest, south-eastern slopes, 750 m, 2.6.1995, F. Özen 1519.

Calystegia silvatica (Kit.) Griseb.

Around Karaköy Village, *Fagus orientalis* forest clearings, northern slopes, 1150 m, 20.7.1994, Özen 1255.

BORAGINACEAE

Myosotis arvensis (L.) Hill subsp. *arvensis*

Around deer production area, *Pinus sylvestris* forest, south-eastern slopes, 1250 m, 7.6.1995, F. Özen 1476, Euro-Siberian element.

M. sicula Guss.

Around deer production area, *Pinus sylvestris* forest clearings, 1300 m, 15.6.1995, F. Özen 1538.

Cynoglossum creticum Mill.

Around deer production area, *Pinus sylvestris* forest clearings, 1200 m, 15.6.1995, F. Özen 1553.

(**)C. germanicum Jacq.

Between Ovacık and deer production area, *Fagus orientalis* forest clearings, north-eastern slopes, 1350 m, 7.6.1995, F. Özen 1486, Euro-Siberian element.

Lithospermum officinale L.

Between Ovacık and deer production area, *Fagus orientalis* forest clearings, 1250 m, 15.6.1995, F. Özen 1608, Euro-Siberian element.

- Echium italicum L.**
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1180 m, 20.7.1994, F. Özen 1315, Mediterranean element?
- E. vulgare L.**
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1200 m, 13.6.1995, F. Özen 1668, Euro-Siberian element.
- Onosma aucheranum DC.**
Around Yukarı Narlı Village, field margins, 500 m, 8.6.1995, F. Özen 1555, East Mediterranean element.
- Anchusa leptophylla Roemer & Schult. subsp. *leptophylla***
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1200 m, 13.6.1995, F. Özen 1671.
- SCROPHULARIACEAE**
Verbascum spectabile M.Bieb. var. *spectabile*
Around Karaköy Village, streamside, 1130 m, 1.8.1995, F. Özen 1461, Euxine element.
Scrophularia scopolii [Hoppe ex] Pers. var. *scopolii*
Around fire observation tower, 1675 m, 1.8.1995, F. Özen 1450.
- S. canina L. subsp. *bicolor* (Sm.) Greuter**
Around Zanbal Village, 1250 m, 12.6.1995, F. Özen 1669, East Mediterranean element.
- Digitalis ferruginea L. subsp. *ferruginea***
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, *Pinus sylvestris* forest clearings, north-western slopes, 1180 m, 15.7.1994, F. Özen 1321, Euro-Siberian element.
- D. ferruginea L. subsp. *schischkinii* (Ivan.) Werner**
Around Gölköy-Asarcık forest store, 1080 m, 13.7.1995, F. Özen 1468, Euxine element.
- Veronica anagallis-aquatica L.**
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, streamsides, 1250 m, 23.7.1994, F. Özen 1310.
- V. chamaedrys L.**
Around deer production area, *Pinus sylvestris-Daphne pontica* association, 1350 m, 15.6.1995, F. Özen 1477, Euro-Siberian element.
- V. orientalis Mill. subsp. *orientalis***
Around Tahtaköprü Village, *Pinus sylvestris* forest clearings, south-western slopes, 1150 m, 10.7.1995, F. Özen 1246, Irano-Turanian element.
- V. multifida L.**
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1250 m, 15.6.1995, F. Özen 1673, Endemic, Irano-Turanian element.
- Melampyrum arvense L. var. *arvense***
Around Göltürkmen Village, *Fagus orientalis* forest clearings, 1110 m, 20.7.1994, F. Özen 1219, Euro-Siberian element.
- Parentucellia viscosa (L.) Caruel**
Around deer production area, *Pinus sylvestris* forest, 1300 m, 23.6.1995, F. Özen 1604, Mediterranean element.
- OROBANCHACEAE**
Orobanche nana Noe ex Beck
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, north-west, 1300 m, 8.6.1995, F. Özen 1493.
- GLOBULARIACEAE**
Globularia trichosantha Fisch. & C.A.Mey.
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1200 m, 8.6.1995, F. Özen 1694.
- VERBENACEAE**
Verbena officinalis L.
Around Karaköy Village, *Pinus sylvestris* forest clearings, 1150 m, 20.7.1994, F. Özen 1242.
- LAMIACEAE (Labiatae)**
Ajuga orientalis L.
Around deer production area, *Pinus sylvestris* forest, 1300 m, 7.6.1995, F. Özen 1537.
- Teucrium chamaedrys L. subsp. *chamaedrys***
Around Mezraa, *Pinus nigra-Juniperus oxycedrus* association, south-eastern slopes, 750 m, 8.6.1995, F. Özen 1524, Euro-Siberian element.
- T. chamaedrys L. subsp. *syspirense* (K.Koch) Rech.f.**
Around Gölalan Village, *Fagus orientalis* forest clearings, southern slopes, 1110 m, 13.7.1994, F. Özen 1211, Irano-Turanian element.
- T. polium L.**
Around Yukarı Narlı Village, field margins, 500 m, 8.6.1995, F. Özen 1563.
- Marrubium astracanicum Jacq. subsp. *astracanicum***
Around deer production area, *Pinus sylvestris-Daphne pontica* association, eastern slopes, 1300 m, 20.7.1994, F. Özen 1351.
- Sideritis montana L. subsp. *montana***
Around Gölalan Village, *Fagus orientalis* forest clearings, eastern slopes, 1150 m, 13.7.1995, F. Özen 1215, Mediterranean element.
- Stachys iberica M.Bieb. subsp. *iberica* var. *iberica***
Around Yukarı Narlı Village, field margins, 500 m, 8.6.1995, F. Özen 1551, Irano-Turanian element.
- S. annua (L.) L. subsp. *annua* var. *annua***
Around deer production area, *Pinus sylvestris* forest, 1350 m, 7.6.1995, F. Özen 1589.
- S. officinalis (L.) Trevis. subsp. *officinalis***
Around Gölalan Village, *Pinus nigra* forest, southern slopes, 1130 m, 31.7.1994, F. Özen 1417, Euro-Siberian element.
- Prunella vulgaris L.**
Around Karaköy Village, *Pinus sylvestris* forest clearings, northern slopes, 1100 m, 13.7.1995, F. Özen 1263, Euro-Siberian element.
- Origanum vulgare L. subsp. *viride* (Boiss.) Hayek**
Around Gölalan Village, *Pinus nigra* forest, southern slopes, 1160 m, 31.7.1994, F. Özen 1416.

| | | |
|---|---|--|
| Calamintha grandiflora (L.) Moench | P. lanceolata L. | E. amygdaloides L. var. amygdaloides |
| Summit of Kunduz Mountain, <i>Fagus orientalis</i> - <i>Galium odoratum</i> association, northern slopes, 1720 m, 1.8.1994, F. Özen 1375, Euro-Siberian element. | Around deer production area, <i>Pinus sylvestris</i> forest clearings, 1250 m, 9.6.1995, F. Özen 1590. | Around Karaköy Village, <i>Fagus orientalis</i> forest, north-western slopes, 1150 m, 11.7.1995, F. Özen 1258, Euro-Siberian element. |
| C. nepeta (L.) Savi subsp. <i>glandulosa</i> (Req.) P.W.Ball | THYMELAEACEAE | URTICACEAE |
| Around Sofular Village, <i>Quercus cerris</i> - <i>Carpinus orientalis</i> association, 1040 m, 2.8.1996, F. Özen 1702. | Daphne pontica L. | Urtica dioica L. |
| Clinopodium vulgare L. subsp. <i>vulgare</i> | Around deer production area, <i>Pinus sylvestris</i> - <i>Daphne pontica</i> association, 1300 m, 7.6.1995, F. Özen 1473, Euxine element. | Around Karaköy Village, <i>Pinus sylvestris</i> forest clearings, 1150 m, 20.7.1994, F. Özen 1241, Euro-Siberian element. |
| Summit of Kunduz Mountain, <i>Fagus orientalis</i> - <i>Galium odoratum</i> association, 1720 m, 1.8.1994, F. Özen 1379. | SANTALACEAE | MORACEAE |
| (**)Thymus longicaulis C.Presl subsp. <i>longicaulis</i> var. <i>subisophyllus</i> (Borbás) Jalas | Thesium divaricatum Jan ex Mert. & Koch | Ficus carica L. subsp. <i>carica</i> |
| Around deer production area, <i>Pinus sylvestris</i> forest, 1300 m, 7.6.1995, F. Özen 1546. | Around Sofular Village, <i>Quercus cerris</i> - <i>Carpinus orientalis</i> association, 1020 m, 1.8.1994, F. Özen 1423, Euro-Siberian element. | Around Sirbaşmak Village, 400 m, 8.6.1995, F. Özen 1613. |
| Mentha longifolia (L.) Hudson subsp. <i>typhoides</i> (Briq.) Harley var. <i>typhoides</i> | ARISTOLOCHIACEAE | JUGLANDACEAE |
| Around Gölalan Village, <i>Fagus orientalis</i> forest clearings, southern slopes, 1110 m, 13.7.1994, F. Özen 1216. | Aristolochia pontica Lam. | (*)Juglans regia L. |
| Salvia viridis L. | Around fire observation tower, <i>Fagus orientalis</i> forest, north-eastern slopes, 1675 m, 31.7.1994, F. Özen 1391, Euxine element. | Around Sirbaşmak Village, 400 m, 8.6.1995, F. Özen 1614. |
| Around Gölköy Village, field margins, 450 m, 1.6.1995, F. Özen 1622, Mediterranean element. | EUPHORBIACEAE | FAGACEAE |
| S. verticillata L. subsp. <i>amasiaca</i> (Freyn & Bornm.) Bornm. | Euphorbia orientalis L. | Fagus orientalis Lipsky |
| Around Gölköy Village, field margins, 500 m, 8.6.1995, F. Özen 1621, Irano-Turanian element? | Around Gölalan Village, <i>Pinus nigra</i> - <i>Juniperus oxycedrus</i> association, eastern slopes, 930 m, 31.7.1994, F. Özen 1430, Irano-Turanian element. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, <i>Fagus orientalis</i> - <i>Galium odoratum</i> association, northern slopes, 1110 m, 13.7.1994, F. Özen 1239, Euro-Siberian element. |
| PLANTAGINACEAE | (**)E. platyphyllos L. | Quercus petraea (Matt.) Liebl. subsp. <i>iberica</i> (Steven ex M.Bieb.) Krassiln. |
| Plantago major L. subsp. <i>major</i> | Around deer production area, <i>Pinus sylvestris</i> - <i>Daphne pontica</i> association, north-eastern slopes, 1350 m, 20.7.1994, F. Özen 1344. | Between Ovacık and Karaköy Villages, 1030 m, 13.7.1994, F. Özen 1366. |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, <i>Pinus sylvestris</i> forest clearings, 1200 m, 9.6.1995, F. Özen 1687. | E. stricta L. | Q. infectoria Olivier subsp. <i>infectoria</i> |
| P. major L. subsp. <i>intermedia</i> (Gilib.) Lange | Around deer production area, <i>Pinus sylvestris</i> - <i>Daphne pontica</i> association, eastern slopes, 1350 m, 20.7.1994, F. Özen 1360, Euro-Siberian element. | Around Yukarı Narlı Village, <i>Pinus nigra</i> - <i>Juniperus oxycedrus</i> association, northern slopes, 200 m, 18.6.1993, F. Özen 1698, Euro-Siberian element. |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1150 m, 23.7.1995, F. Özen 1295. | E. helioscopia L. | Q. cerris L. var. <i>cerris</i> |
| P. holosteum Scop. | Around Tahtaköprü Village, <i>Pinus sylvestris</i> forest clearings, western slopes, 1375 m, 15.6.1995, F. Özen 1598. | Around Mezraa, <i>Quercus cerris</i> - <i>Carpinus orientalis</i> association, southern slopes, 550 m, 13.7.1994, F. Özen 1324, Mediterranean element. |
| Around fire observation tower, <i>Fagus orientalis</i> forest clearings, eastern slopes, 1675 m, 1.8.1995, F. Özen 1456, Mediterranean element. | E. falcata L. subsp. <i>falcata</i> var. <i>falcata</i> | CORYLACEAE |
| | Around Sofular Village, <i>Pinus nigra</i> - <i>Juniperus oxycedrus</i> association, 920 m, 3.8.1995, F. Özen 1432. | Carpinus orientalis Mill. subsp. <i>orientalis</i> |
| | | Around Göltürkmen Village, <i>Quercus cerris</i> - <i>Carpinus orientalis</i> association, 1040 m, 31.7.1994, F. Özen 1697. |

- C. betulus L.**
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, *Fagus orientalis* forest, 1050 m, 9.6.1995, F. Özen 1643, Euro-Siberian element.
- Corylus avellana L. var. *avellana***
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, streamside, north-eastern slopes, 1130 m, 29.7.1994, F. Özen 1442, Euro-Siberian element.
- SALICACEAE**
Salix alba L.
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, streamside, 1200 m, 23.7.1994, F. Özen 1320, Euro-Siberian element.
- RUBIACEAE**
Asperula involucrata Wahlenb.
Around Gölalan Village, *Quercus petraea* forest, 950 m, 8.6.1995, F. Özen 1498, Euxine element.
- A. arvensis L.**
Around deer production area, *Pinus sylvestris* forest clearings, south-eastern slopes, 1300 m, 15.6.1995, F. Özen 1545, Mediterranean element.
- Galium rotundifolium L.**
Between Ovacık and deer production area, *Fagus orientalis* forest, north-eastern slopes, 1500 m, 8.8.1995, F. Özen 1403, Euro-Siberian element.
- G. odoratum (L.) Scop.**
Around Öğürlü Village, *Quercus cerris-Carpinus orientalis* association, south-eastern slopes, 1000 m, 1.6.1995, F. Özen 1501, Euro-Siberian element.
- (**)G. elongatum C.Presl**
Around Gölköy Village, field margins, 550 m, 1.6.1995, F. Özen 1623, Euro-Siberian element.
- G. verum L. subsp. *verum***
Around Sofular Village, *Fagus orientalis* forest clearings, 1100 m, 10.7.1995, F. Özen 1245, Euro-Siberian element.
- G. verum L. subsp. *glabrescens* Ehrend.**
Around Gölalan Village, *Fagus orientalis* forest clearings, 1110 m, 20.7.1994, F. Özen 1199, Irano-Turanian element?
- G. paschale Forssk.**
Around Öğürlü Village, *Quercus cerris-Carpinus orientalis* association, north-eastern slopes, 1000 m, 1.6.1995, F. Özen 1500, East Mediterranean (mountain) element.
- G. divaricatum Pourr. ex Lam.**
Around Gölalan Village, *Pinus nigra* forest, 1130 m, 23.6.1994, F. Özen 1413, Mediterranean element.
- (**)G. peplidifolium Boiss.**
Around deer production area, *Pinus sylvestris* forest, south-eastern slopes, 1400 m, 9.6.1994, F. Özen 1575, East Mediterranean element.
- MONOCOTYLEDONAE**
ARACEAE
()Arum italicum Mill.**
Between Ovacık and Karaköy Villages, *Fagus orientalis* forest clearings, 1110 m, 20.7.1994, F. Özen 1204.
- LILIACEAE**
Polygonatum multiflorum (L.) All.
Around deer production area, *Fagus orientalis* forest, 1300 m, 8.6.1995, F. Özen 1645.
- Scilla autumnalis L.**
Around Sofular Village, *Quercus cerris-Carpinus orientalis* association, northern slopes, 1040 m, 29.7.1994, F. Özen 1420, Mediterranean element.
- Muscari armeniacum Leichtlin ex Baker**
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, *Pinus sylvestris* forest clearings, 1200 m, 8.6.1995, F. Özen 1663.
- M. neglectum Guss.**
Around Yukarı Narlı Village, *Pinus nigra* forest, south-western slopes, 350 m, 2.5.1993, F. Özen 1192.
- ORCHIDACEAE**
Neottia nidus-avis (L.) J.M.C.Rich.
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, *Fagus orientalis-Galium odoratum* association, northern slopes, 1250 m, 3.8.1996, F. Özen 1708, Euro-Siberian element.
- Cephalanthera rubra (L.) J.M.C.Rich.**
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 900 m, 8.6.1995, F. Özen 1505.
- C. damasonium (Mill.) Druce**
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, Northeast, 1000 m, 8.6.1995, F. Özen 1497, Euro-Siberian element.
- Epipactis pontica Taub.**
Around fire observation tower, 1670 m, 31.7.1994, F. Özen 1397, Endemic, Euxine element.
- Epipogium aphyllum Sw.**
Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, east, 1200 m, 20.7.1994, F. Özen 1333, Euro-Siberian element.
- Platanthera chlorantha (Custer) Rchb.**
Around deer production area, *Pinus sylvestris-Daphne pontica* association, eastern slopes, 1400 m, 3.8.1996, F. Özen 1706.
- JUNCACEAE**
Luzula pseudosudetica (V.I.Krecz.) V.I.Krecz. apud V.I.Krecz. & Gontsch.
Around Soruk Village, *Pinus sylvestris-Daphne pontica* association, south-eastern slopes, 1300 m, 20.6.1995, F. Özen 1480, Euxine (mountain) element.
- L. multiflora (Ehrh. ex Retz.) Lej.**
Around Karaköy Village, *Pinus sylvestris* forest, 1375 m, 7.6.1995, F. Özen 1583.
- CYPERACEAE**
()Carex pendula Huds.**
Around Soruk Village, streamside, 1000 m, 1.6.1995, F. Özen 1639, Euro-Siberian element.
- C. sylvatica Huds. subsp. *sylvatica***
Around Soruk Village, *Pinus sylvestris* forest, 1000 m, 1.6.1995, F. Özen 1638.
- C. flacca Schreber subsp. *serrulata* (Biv.) Greuter**
Around deer production area, *Pinus sylvestris* forest, 1350 m, 7.6.1995, F. Özen 1600, Mediterranean element.
- (**)C. acuta L.**
Around Gölalan Village, *Pinus nigra* forest, 1130 m, 31.7.1994, F. Özen 1408, Euro-Siberian element.

| | | |
|--|---|--|
| POACEAE (Gramineae) | (**) <i>Avena fatua</i> L. var. <i>glabrata</i> | <i>Dactylis glomerata</i> L. subsp. <i>glomerata</i> |
| <i>Aegilops geniculata</i> Roth | Peterm. | Around Soruk Village, <i>Fagus orientalis</i> - <i>Galium odoratum</i> association, 1720 m, 31.7.1994, F. Özen 1383, Euro-Siberian element. |
| Around Yukarı Narlı Village, <i>Pinus nigra</i> - <i>Juniperus oxycedrus</i> association, 300 m, 18.6.1993, F. Özen 1699, Mediterranean element. | Around Gölalan Village, <i>Fagus orientalis</i> forest clearings, 1100 m, 20.7.1994, F. Özen 1232, Euro-Siberian element. | <i>D. glomerata</i> L. subsp. <i>hispanica</i> (Roth) Nyman |
| <i>Triticum aestivum</i> L. | <i>Koeleria cristata</i> (L.) Pers. | Around Mezraa, <i>Pinus nigra</i> - <i>Juniperus oxycedrus</i> association, south-eastern slopes, 750 m, 8.6.1995, F. Özen 1529. |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1150 m, 20.7.1994, F. Özen 1304. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1180 m, 20.7.1994, F. Özen 1301. | <i>Cynosurus cristatus</i> L. |
| <i>Hordeum murinum</i> L. subsp. <i>glaucum</i> (Steud.) Tzvelev | <i>Festuca heterophylla</i> Lam. | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, field margins, 1100 m, 20.7.1994, F. Özen 1308, Euro-Siberian element. |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1250 m, 8.6.1995, F. Özen 1684. | Around deer production area, <i>Pinus sylvestris</i> forest clearings, south-western slopes, 1300 m, 8.6.1995, F. Özen 1649, Euro-Siberian element. | <i>C. echinatus</i> L. |
| <i>H. bulbosum</i> L. | <i>Lolium rigidum</i> Gaudin var. <i>rigidum</i> | Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, ponds, 1200 m, 7.6.1995, F. Özen 1532, Mediterranean element. |
| Around deer production area, 1350 m, 7.6.1995, F. Özen 1586. | Around Gölalan Village, <i>Fagus orientalis</i> forest clearings, north-eastern slopes, 1110 m, 20.7.1994, F. Özen 1235. | <i>Briza media</i> L. |
| <i>Bromus tectorum</i> L. | <i>Poa alpina</i> L. subsp. <i>fallax</i> F.Herm. | Around Yukarı Narlı Village, <i>Pinus nigra</i> ssp. <i>pallasiana</i> - <i>Juniperus oxycedrus</i> association, 850 m, 31.7.1994, F. Özen 1433. |
| Ovacık Village, around Burhan Zorlu Education Centre belonging to the Ministry of Forestry, 1200 m, 8.6.1995, F. Özen 1677. | Around deer production area, <i>Pinus sylvestris</i> - <i>Daphne pontica</i> association, south-eastern slopes, 1300 m, 7.6.1995, F. Özen 1479. | |

Vegetation

There are two vegetation types in the study area: forest and degraded forests. Vegetation is stratified longitudinally. It is known that altitude, direction, topography, temperature and precipitation play an important role in the stratification (Özen & Kilinç, 1995). In this study it was found that *Quercus cerris* var. *cerris* and *Carpinus orientalis* were generally dominant and *Pinus nigra* subsp. *pallasiana* was sometimes dominant at low altitudes in the region. The dominance of *Pinus nigra* subsp. *pallasiana* increases with altitude up to 900 m. After 900 m *Quercus cerris* var. *cerris*, *Carpinus orientalis* and *Pinus nigra* subsp. *pallasiana* are almost equally dominant. *Pinus sylvestris* is dominant especially on southern slopes, whereas *Fagus orientalis* is dominant especially on northern slopes. Both are spread together at the higher vegetation stratum. *Fagus orientalis* is only dominant at the highest vegetation stratum in the study area.

A summary of vegetation stratification in the study area is given below:

400-600 m *Quercus cerris*-*Carpinus orientalis*-*Pinus nigra*

| | |
|-------------|--|
| 600-1150 m | <i>Pinus nigra</i> - <i>Quercus cerris</i> |
| 900-1050 m | <i>Quercus cerris</i> - <i>Carpinus orientalis</i> |
| 1000-1500 m | <i>Pinus sylvestris</i> - <i>Fagus orientalis</i> |
| 1450-1740 m | <i>Fagus orientalis</i> |

The associations found in the study area are as follows:

Degraded forest vegetation

1- *Carpino-Quercetum cerridis* Kutbay & Kilinç 1995 (Table 1)

Characteristic and differential species of this association are *Quercus cerris* var. *cerris*, *Carpinus orientalis* subsp. *orientalis* and *Quercus petraea* subsp. *iberica*. *Carpino-Quercetum cerridis* occurs between 400 and 1040 m. It is spread on brown forest soils and exhibits three vegetation layers, namely tree, shrub and herb. The total coverage of the tree layer varies between 5 and 90% and 5 and 10 m in height. The most common species in the tree layer are *Pinus nigra* subsp. *pallasiana* and *Fagus orientalis*. The total coverage and height of the shrub layer are 30 and 90% and 1 and 3 m, respectively. The most common species in the shrub layer are *Quercus*

Table 1. Carpino-Quercetum cerridis Kutbay & Kılıç 1995

| Quadrat No | 23 | 24 | 46 | 47 | 80 | 81 | 96 | 72 | 73 | |
|---|-----|-----|------|------|------|------|------|------|------|----------|
| Altitude (m) | 600 | 400 | 1040 | 1020 | 1000 | 1020 | 1000 | 1000 | 1000 | |
| Inclination (%) | - | - | - | 10 | - | - | 10 | 25 | - | |
| Exposure | - | - | - | SW | - | - | E | NE | - | |
| Height of the tree layer (m) | 5 | 10 | 7 | 10 | 10 | 8 | 8 | 5 | 5 | |
| Coverage of the tree layer (%) | - | - | 90 | 90 | 10 | 5 | - | 80 | 90 | |
| Height of the shrub layer (m) | - | 3 | 1.5 | 1 | 5 | 2 | 2 | 2 | 1 | |
| Coverage of the shrub layer (%) | 80 | 80 | 30 | 30 | 90 | 90 | 80 | 50 | 70 | |
| Height of the herb layer (cm) | 10 | 10 | 15 | 15 | 15 | 10 | 10 | 15 | 15 | |
| Coverage of the herb layer (%) | 15 | 15 | 20 | 15 | 10 | 10 | 10 | 20 | 30 | Presence |
| Characteristic and differential species of the association | | | | | | | | | | |
| <i>Quercus cerris</i> var. <i>cerris</i> | 44 | 33 | 45 | 44 | 44 | 45 | 33 | 12 | 44 | V |
| <i>Carpinus orientalis</i> subsp. <i>orientalis</i> | 23 | 22 | 33 | 33 | 22 | 33 | 33 | 22 | 22 | V |
| <i>Quercus petraea</i> subsp. <i>iberica</i> | 11 | 12 | 22 | 22 | 22 | 22 | 33 | 44 | 11 | V |
| Characteristic species of Querco-Carpinetalia orientalis and Carpino-Acerion (*) | | | | | | | | | | |
| <i>Crataegus monogyna</i> subsp. <i>monogyna</i> | 12 | . | 11 | 11 | 11 | 22 | +2 | . | . | IV |
| <i>Lathyrus laxiflorus</i> ssp. <i>laxiflorus</i> | +1 | +1 | . | 11 | 11 | . | . | +2 | 11 | IV |
| <i>Dorycnium pentaphyllum</i> subsp. <i>anatolicum</i> | . | . | . | +2 | +1 | . | . | +2 | . | II |
| (*) <i>Cyclamen coum</i> var. <i>coum</i> | . | 12 | . | . | . | +1 | . | . | +1 | II |
| (*) <i>Asperula involucrata</i> | . | 11 | . | . | . | . | . | . | +2 | II |
| Characteristic species of Quercetea pubescens | | | | | | | | | | |
| <i>Pinus nigra</i> subsp. <i>pallasiana</i> | . | 11 | +1 | +1 | +1 | 11 | . | +1 | +1 | IV |
| <i>Sorbus torminalis</i> var. <i>pinnatifida</i> | . | . | . | +1 | +1 | . | +1 | +1 | 11 | III |
| <i>Juniperus oxycedrus</i> subsp. <i>oxycedrus</i> | . | . | +2 | +2 | . | 11 | . | . | . | II |
| <i>Acer campestre</i> subsp. <i>leiocarpum</i> | . | . | . | +1 | . | . | . | . | . | I |
| Characteristic species of Rhododendro-Fagetalia orientalis | | | | | | | | | | |
| <i>Fagus orientalis</i> | . | . | +1 | . | +1 | +1 | +1 | . | . | II |
| <i>Daphne pontica</i> | . | +1 | +1 | . | +1 | . | . | . | . | II |
| <i>Rubus hirtus</i> | . | . | . | . | . | 12 | . | . | . | I |
| Characteristic species of Querco-Fagea and Quero-Fagetea (*) | | | | | | | | | | |
| (*) <i>Euphorbia amygdaloides</i> var. <i>amygdaloides</i> | . | . | +1 | +1 | . | 11 | . | +1 | +1 | III |
| <i>Primula vulgaris</i> subsp. <i>vulgaris</i> | +1 | . | +1 | . | . | . | . | 11 | 11 | III |
| <i>Fragaria vesca</i> | . | . | 11 | +1 | . | . | . | . | +2 | II |
| <i>Clinopodium vulgare</i> subsp. <i>vulgare</i> | . | +1 | . | . | . | . | . | . | . | I |
| <i>Teucrium chamaedrys</i> subsp. <i>chamaedrys</i> | . | . | . | . | . | . | +2 | . | . | I |
| <i>Geum urbanum</i> | . | . | . | . | . | . | . | . | +1 | I |
| <i>Veronica chamaedrys</i> | . | . | . | . | . | . | . | . | +1 | I |
| Other Species | | | | | | | | | | |
| <i>Rosa canina</i> | 11 | +1 | 11 | 11 | +1 | . | +1 | +2 | +2 | V |
| <i>Cornus sanguinea</i> subsp. <i>orientalis</i> | +1 | . | +1 | . | +1 | . | . | . | . | II |
| <i>Genista lydia</i> var. <i>lydia</i> | . | +2 | . | . | . | . | . | +1 | . | II |
| <i>Crataegus angustifolia</i> | . | . | . | . | . | . | . | 22 | 22 | II |
| <i>Bellis perennis</i> | +1 | . | . | . | . | +1 | . | . | . | I |
| <i>Silene vulgaris</i> var. <i>vulgaris</i> | . | +1 | . | . | . | . | . | . | . | I |
| <i>Calamintha nepeta</i> | . | . | +1 | . | . | . | . | . | . | I |
| <i>Scilla autumnalis</i> | . | . | +1 | . | . | . | . | . | . | I |
| <i>Pilosella hoppeana</i> subsp. <i>testimonialis</i> | . | . | +1 | . | . | . | . | . | . | I |
| <i>Thesium divaricatum</i> | . | . | . | +2 | . | . | . | . | . | I |
| <i>Plantago lanceolata</i> | . | . | . | +1 | . | . | . | . | . | I |
| <i>Helianthemum nummularium</i> subsp. <i>nummularium</i> | . | . | . | . | . | . | +1 | . | . | I |
| <i>Luzula multiflora</i> | . | . | . | . | . | . | . | +1 | . | I |
| <i>Vicia crocea</i> | . | . | . | . | . | . | . | +1 | . | I |
| <i>Cephalanthera damasonium</i> | . | . | . | . | . | . | . | 11 | . | I |
| <i>Galium odoratum</i> | . | . | . | . | . | . | . | +1 | . | I |
| <i>Galium paschale</i> | . | . | . | . | . | . | . | +1 | . | I |

cerris var. *cerris*, *Carpinus orientalis*, *Quercus petraea* subsp. *iberica*, *Crataegus monogyna* subsp. *monogyna* and *Rosa canina*. The total coverage of the herb layer varies between 10 and 30% and 10 and 15 cm in height. *Lathyrus laxiflorus* subsp. *laxiflorus*, *Euphorbia amygdaloides* var. *amygdaloides* and *Primula vulgaris* subsp. *vulgaris* are common species in this layer.

Carpino-Quercetum cerridis is more degraded than the other associations in the study area. This association includes various species belonging to various syntaxa units. In particular, species belonging to the class QUERCETEA PUBESCENTIS (Oberd. 1948) Doing Kraft 1955, the order QUERCO-CARPINETALIA ORIENTALIS Quezel, Barbero et Akman 1980 and the alliance CARPINO-ACERION Quezel, Barbero et Akman 1978 are richer than the other syntaxonomic units. The class QUERCETEA PUBESCENTIS is characterised by *Quercus cerris* var. *cerris*, *Pinus nigra* subsp. *pallasiana*, *Sorbus torminalis* var. *pinnatifida*, *Juniperus oxycedrus* subsp. *oxycedrus* and *Acer campestre* subsp. *leiocarpum*. The order QUERCO-CARPINETALIA ORIENTALIS and the alliance CARPINO-ACERION are represented by *Carpinus orientalis* subsp. *orientalis*, *Quercus petraea* subsp. *iberica*, *Crataegus monogyna* subsp. *monogyna*, *Lathyrus laxiflorus* subsp. *laxiflorus*, *Cyclamen coum* var. *coum* and *Asperula involucrata*. On the basis of the information given above, it is apparent that this association should be included in the order QUERCO-CARPINETALIA ORIENTALIS.

Forest vegetation

2- *Junipero-Pinetum nigrae* Özen & Kilinç as. nov.
(Table 2)

This association (holotype table 2, quadrat number 25) is widespread on the southern and eastern slopes between 400 and 1130 m and on brown forest soils in the study area. The characteristic and differential species of the association are *Pinus nigra* subsp. *pallasiana*, *Juniperus oxycedrus* subsp. *oxycedrus* and *Dorycnium pentaphyllum* subsp. *anatolicum*. This association comprises tree, shrub and herb layers. The tree layer is represented by *Pinus nigra* subsp. *pallasiana* and *Fagus orientalis*. The total coverage and height vary between 75 and 100% and 10 and 15 m in the tree layer, respectively.

The shrub layer is between 0.5 and 2 m in height and total coverage varies between 10 and 70%. The most

common species in this layer are *Juniperus oxycedrus* subsp. *oxycedrus*, *Quercus cerris* var. *cerris* and *Carpinus orientalis* subsp. *orientalis*. The total coverage and height of the herb layer are between 10 and 40% and 5 and 20 cm, respectively. The dominant species in the herb layer are *Dorycnium pentaphyllum* subsp. *anatolicum*, *Dorycnium graecum*, *Dactylis glomerata* subsp. *hispanica* and *Polygonum pruinosa*.

Characteristic species of different syntaxonomic units are found together in the association *Junipero-Pinetum nigrae*. The order RHODODENDRO-FAGETALIA ORIENTALIS Quezel, Barbero et Akman 1980 is characterised by *Daphne pontica* and *Fagus orientalis*. The class QUERCETEA ILCIS Braun-Blanquet 1931 is only represented by *Phillyrea latifolia*. The frequency and cover percentage of species in this order and class are very low. The components of CISTO-MICROMERIETEA Raus 1979 are *Pteridium aquilinum*, *Teucrium polium*, *Origanum vulgare* subsp. *viride*, *Psoralea bituminosa* and *Cistus creticus*. The spread of taxa belonging to this class shows, to some extent, some human activity on the association. Species belonging to the class QUERCETEA PUBESCENTIS and the order QUERCO-CARPINETALIA ORIENTALIS are dominant in the association *Junipero-Pinetum nigrae*. Therefore, it is more appropriate to include this association in those syntaxa units.

3- *Daphno-Pinetum sylvestris* Kutbay & Kilinç 1995
(Table 3)

The characteristic and differential species of this association are *Pinus sylvestris*, *Daphne pontica* and *Galium rotundifolium*. *Daphno-Pinetum sylvestris* is widespread, especially on the southern and eastern slopes between 1000 and 1400 m and on grey-brown podsolic soils in the study area.

The association comprises tree, shrub and herb layers. The tree layer is represented by *Pinus sylvestris* and *Fagus orientalis* having total coverage between 80 and 100% and 5 and 30 m height, respectively. The total coverage and height of the shrub layer vary between 15 and 35% and 0.5 and 1 m, respectively. The dominant species of this layer are *Daphne pontica*, *Pyracantha coccinea* and *Rubus hirtus*. The total coverage and height of the herb layer vary between 10 and 30% and 10 and 30 cm, respectively. The most common species are *Galium rotundifolium*, *Euphorbia amygdaloides* var. *amygdaloides*, *Sanicula europaea*, *Primula vulgaris* subsp. *vulgaris* and *Fragaria vesca* in this layer.

Table 2. Juniper-Pinetum nigrae Özen & Kılıç as. nov. ** Type: Quadrat 25

Euphorbia stricta subsp. *farcta* var. *farcta* (85). Campanula rapunculus subsp. *rapunculus* (44), *calum divaricatum* (44). Cynosurus echinatus (44). Gentiana lutea var. *caneensis* (76). Linum suffruticosum (76). Convolvulus cantabrica (76). Inula viscosa subsp. *sibirica* (76). Filipendula vulgaris (76). Urtica dioica subsp. *canescens* (76). Anemone nemorosa subsp. *canescens* (76). Convallaria majalis (76). Lathyrus aphaca var. *paeoniflora* (76). Filago aegyptiaca (76). Artea squamata (62).

Table 3. Daphno-Pinetum sylvestris Kutluay & Kılımcı 1995

| | Quadrat No | 20 | 21 | 82 | 1 | 2 | 6 | 66 | 67 | 99 | 77 | 78 | 12 | 13 | 63 | 64 |
|--|-----------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| | Altitude (m) | 1350 | 1400 | 1300 | 1200 | 1250 | 1350 | 1300 | 1000 | 1350 | 1330 | 1350 | 1300 | 1350 | 1200 | 1200 |
| | Inclination (%) | 25 | 10 | 15 | 30 | 20 | 30 | 40 | 50 | 15 | 10 | 25 | 35 | 25 | 20 | 20 |
| Exposure | E | SE | E | SE | E | NE | N | SE | SW | E | NE | - | - | SE | - | - |
| Height of the tree layer (m) | 15 | 15 | 25 | 30 | 25 | 25 | 20 | 20 | 20 | 30 | 20 | 20 | 25 | 25 | 25 | 25 |
| Coverage of the tree layer (%) | 85 | 90 | 100 | 95 | 90 | 90 | 80 | 95 | 95 | 100 | 90 | 90 | 100 | 90 | 90 | 90 |
| Height of the shrub layer (m) | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 1 | 0.5 | 1 | 0.75 | 0.5 | 0.5 | 0.75 | 0.5 | 0.5 | 0.5 |
| Coverage of the shrub layer (%) | 15 | 30 | 20 | 20 | 25 | 30 | 25 | 30 | 35 | 25 | 25 | 15 | 30 | 25 | 30 | 20 |
| Height of the herb layer (cm) | 10 | 20 | 10 | 25 | 30 | 30 | 25 | 30 | 25 | 30 | 15 | 25 | 20 | 25 | 20 | 25 |
| Coverage of the herb layer (%) | 10 | 30 | 10 | 25 | 30 | 30 | 25 | 30 | 25 | 30 | 15 | 25 | 20 | 25 | 20 | 25 |
| Characteristic and differential species of the association | | | | | | | | | | | | | | | | |
| <i>Pinus sylvestris</i> | 45 | 45 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 55 | 44 | 44 |
| <i>Daphne pontica</i> | 22 | 22 | +1 | 11 | 22 | 12 | 22 | 22 | +1 | 11 | +1 | 22 | 22 | 22 | 22 | 22 |
| <i>Gallium rotundifolium</i> | | | | | | | | | | | | | | | +1 | IV |
| Characteristic species of Querco-Fagetea | | | | | | | | | | | | | | | | |
| <i>Euphorbia amygdaloides</i> var. <i>amygdaloides</i> | 12 | 12 | 12 | 11 | 11 | 11 | 22 | 12 | 12 | 22 | 11 | 22 | 22 | 12 | V | V |
| <i>Viola odorata</i> | | | | +1 | | +1 | | | | | +1 | | | | | |
| Stellaria holostea | | | | | | | | | | | | | | | | |
| Characteristic species of Rhododendro-Fageta orientalis | | | | | | | | | | | | | | | | |
| <i>Fagus orientalis</i> | | | +1 | | +1 | 11 | +1 | 11 | +1 | +1 | +1 | +1 | +1 | +1 | | |
| <i>Rubus hirtus</i> | | | | | | +2 | 11 | +1 | +1 | +1 | +1 | +1 | +1 | +1 | | |
| Characteristic species of Querco-Carpinetalia orientalis and Carpino-Acerion (*) | | | | | | | | | | | | | | | | |
| <i>(*) Pyracantha coccinea</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | +1 | |
| <i>(*) Cyclamen coum</i> var. <i>coum</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | +1 | |
| <i>Lathyrus laxiflorus</i> subsp. <i>laxiflorus</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | +2 | |
| <i>Doronicum orientale</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | 22 | |
| <i>Digitalis ferruginea</i> subsp. <i>ferruginea</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | | |
| <i>Geranium asphodeloides</i> subsp. <i>asphodeloides</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | | |
| <i>Crataegus monogyna</i> subsp. <i>monogyna</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | | |
| <i>Dorycnium graecum</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | | |
| Characteristic species of Fageta sylvaticae | | | | | | | | | | | | | | | | |
| <i>Sanicula europaea</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | +1 | V |
| <i>Calamintha grandiflora</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | | |
| <i>Mycelis muralis</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | | |
| <i>Gallium odoratum</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | | |
| Characteristic species of Quercetalia pubescantis | | | | | | | | | | | | | | | | |
| <i>Juniperus oxycedrus</i> subsp. <i>oxycedrus</i> | | | +2 | | | | | | | | | | | | | |
| <i>Comus mas</i> | | | | | | | | | | | | | | | | |
| Characteristic species of Querco-Fageta | | | | | | | | | | | | | | | | |
| <i>Primula vulgaris</i> subsp. <i>vulgaris</i> | 11 | 22 | 11 | +1 | +1 | +1 | 11 | 11 | +1 | 11 | +1 | 11 | 11 | 22 | 11 | V |
| <i>Fragaria vesca</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | +1 | |
| <i>Veronica chamaedrys</i> | | | +1 | +1 | +1 | +1 | | | | | | | | | +1 | |
| <i>Hedera helix</i> | | | +1 | | | | | | | | | | | | | |
| Other Species | | | | | | | | | | | | | | | | |
| <i>Pteridium aquilinum</i> | 12 | 33 | +1 | 33 | 33 | 33 | 33 | 33 | 22 | 11 | 23 | | | 22 | 12 | V |
| <i>Myosotis arvensis</i> var. <i>arvensis</i> | +1 | +2 | | | +2 | | | | | | | | | | | |
| <i>Sambucus ebulus</i> | | | | | | | | | | | | | | | | |
| <i>Lathyrus laxiflorus</i> subsp. <i>angustifolius</i> | | | | | | | | | | | | | | | | |
| <i>Digitalis ferruginea</i> subsp. <i>schischkinii</i> | | | | | | | | | | | | | | | | |
| <i>Phosella hoppeana</i> subsp. <i>testimoniensis</i> | | | | | | | | | | | | | | | | |
| <i>Euphorbia stricta</i> | | | | | | | | | | | | | | | | |
| <i>Prunella vulgaris</i> | 11 | +1 | | | | | | | | | | | | | | |
| <i>Bells perennis</i> | | | | | | | | | | | | | | | | |
| <i>Pilosella x auriculoides</i> | | | | | | | | | | | | | | | | |
| <i>Poa alpina</i> subsp. <i>fallax</i> | | | | | | | | | | | | | | | | |
| <i>Rubus canescens</i> var. <i>canescens</i> | | | | | | | | | | | | | | | | |
| <i>Luzula pseudosudetica</i> | | | | | | | | | | | | | | | | |
| <i>Carex acuta</i> | | | | | | | | | | | | | | | | |
| <i>Cirsium pseudopersonata</i> subsp. <i>pseudopersonata</i> | +1 | | | | | | | | | | | | | | | |
| <i>Gallium peplidifolium</i> | +1 | | | | | | | | | | | | | | | |
| The species of single frequency: <i>Platianthera chlorantha</i> (21), <i>Euphorbia platyphyllus</i> (21), <i>Epipactis apylloides</i> (1), <i>Sambucus nigra</i> (78), <i>Lotus angustissimus</i> (6), <i>Polyistichum setiferum</i> (67), <i>Epiptactis pontica</i> (99), <i>Sedum pallidum</i> var. <i>pallidum</i> (77), <i>Sanguisorba minor</i> subsp. <i>muricata</i> (77), <i>Crataegus tanacetifolia</i> (77), <i>Plantago lanceolata</i> (78), <i>Marrubium astracanicum</i> (78), <i>Rosa canina</i> (63). | | | | | | | | | | | | | | | | |

Daphno-Pinetum sylvestris includes various species belonging to various syntaxa units. The class QUERCO-FAGETEA (Br.-Bl. et Vliegler 1937) Fuk. et Fab. 1968 is represented by *Pinus sylvestris*, *Galium rotundifolium*, *Euphorbia amygdaloides* var. *amygdaloides*, *Viola odorata* and *Stellaria holostea*. The order RHODODENDRO-FAGETALIA ORIENTALIS is characterised by *Fagus orientalis*, *Daphne pontica* and *Rubus hirtus*. The important components of the order QUERCO-CARPINETALIA ORIENTALIS and alliance CARPINO-ACERION are *Pyracantha coccinea*, *Cyclamen coum* var. *coum*, *Lathyrus laxiflorus* subsp. *laxiflorus* and *Doronicum orientale*. The order FAGETALIA SYLVATICA is represented by *Sanicula europaea* and *Calamintha grandiflora*. The class QUERCETEA PUBESCENTIS is characterised by *Juniperus oxycedrus* subsp. *oxycedrus* and *Cornus mas*. The association *Daphno-Pinetum sylvestris* was included in the class QUERCO-FAGETEA and the order RHODODENDRO-FAGETALIA ORIENTALIS when the floristic composition was examined.

4- Galio-Fagetum orientalis Özén & Kılıç as. nov. (Table 4)

This association (holotype table 4, quadrat number 27) is scattered, especially on northern slopes between 1150 and 1740 m and on grey-brown podsolic soils in the study area. The characteristic and differential species of the association are *Fagus orientalis*, *Galium odoratum* and *Rubus hirtus*. The association exhibits tree, shrub and herb layers. The total coverage and height of the tree layer vary between 90 and 100% and 15 and 25 m, respectively. This layer is characterised by *Fagus orientalis* and *Pinus sylvestris*. The total coverage and height of the shrub layer vary between 10 and 20% and 0.5 and 1 m, respectively. The dominant species of the bush layer are *Rubus hirtus* and *Daphne pontica*. The total coverage and height of the herb layer vary between 5 and 40% and 5 and 20 cm, respectively. The most common species in the herb layer are *Galium odoratum*, *Lathyrus laxiflorus* subsp. *laxiflorus*, *Fragaria vesca* and *Urtica dioica*.

The floristic composition of the association includes the characteristic species of the orders FAGETALIA SYLVATICA, RHODODENDRO-FAGETALIA ORIENTALIS and QUERCO-CARPINETALIA ORIENTALIS and the classes QUERCETEA PUBESCENTIS and QUERCO-FAGETEA. The characteristic species of the order FAGETALIA SYLVATICA are *Monotropa hypopitys*,

Calamintha grandiflora, *Sanicula europaea*, and *Neottia nidus-avis*. The important components of the order QUERCO-CARPINETALIA ORIENTALIS and the alliance CARPINO-ACERION are *Lathyrus laxiflorus* subsp. *laxiflorus*, *Cyclamen coum* var. *coum* and *Doronicum orientale*. The class QUERCETEA PUBESCENTIS is only represented by *Cephalanthera rubra*.

From this point of view, the syntaxonomic explanation of the *Galio-Fagetum orientalis* is not easy. However, all characteristic and differential species of this association are typical for the order RHODODENDRO-FAGETALIA ORIENTALIS. Moreover, the class QUERCO-FAGETEA is also well represented in this association. Therefore, it should be considered in the syntaxa cited above.

Conclusions and Comments

From the 306 taxa found in the study area, *Ailanthus altissima*, *Pistacia vera*, *Pyrus communis* subsp. *sativa* and *Juglans regia* are culture plants and 25 taxa are new floristic records for the A5 square in the Flora of Turkey.

The distribution of taxa according to life forms and living periods is shown in Tables 5 and 6, respectively.

As seen in Table 5, the percentage of hemicryptophytes is higher than those of the other life forms. The proportions of therophytes and chamaephytes appear to be remarkably higher than those of the other life form categories (i.e. macrophanerophytes, nanophanerophytes and cryptophytes).

The distribution of taxa in different floristic regions is given in Table 7.

As seen in Table 7, Euro-Siberian elements are more common than the other floristic elements in the research area. This is due to the fact that the research area is situated in the Black Sea region. The lower altitudes of the research area have a Mediterranean microclimate because of its being near Kızılırmak Valley. Therefore, the percentage of Mediterranean elements is high at lower altitudes. In addition, some Irano-Turanian elements are also present, because the study area is located between the Black Sea and Central Anatolian regions.

Table 8 shows how similar the study and nearby areas are in terms of floristic composition.

Özen & Kılıç's study (1995) describes floristic properties the most similar to those in the study area (Table 8). The proportion of endemism in the area is very

Table 4. Galio-Fagetum orientalis Özen & Kılınç as. nov. ** Type: Quadrat 27

The species of single frequency: *Epigaea apifolia* (4), *Cirsium pseudoperssonata* (5), *Sedum pallidum* var. *pallidum* (27), *Seneio vernalis* (27), *Cystopteris fragilis* (29), *Stenocarpus linearis* (31), *Leptosiphon speciosus*-*vemens* (31), *Hypoxis* (31), *Thlaspi* (31), *Campanula rotundifolia* subsp. *hispanica* (31), *Cynoglossum germanicum* (68), *Epilobium lanceolatum* (68), *Pilosella x auriculata* (68), *Sanguisorba minor* subsp. *muraria* (71), *Tussilago farfara* (71), *Crataegus angustifolia* (70), *Rumex acetosa* (38), *Trifolium pratense* (31), *Agrostis capillaris* (31).

Table 5. The distribution of taxa according to life forms.

| Life form | Number of Species | Percentage |
|-------------------|-------------------|------------|
| Macrophanerophyte | 17 | 5.55 |
| Nanophanerophyte | 29 | 9.48 |
| Chamaephyte | 65 | 21.24 |
| Hemicryptophyte | 122 | 39.87 |
| Therophyte | 67 | 21.90 |
| Cryptophyte | 6 | 1.96 |
| Total | 306 | 100.00 |

Table 6. The distribution of taxa according to living periods

| Living period | Number of Species | Percentage |
|---------------|-------------------|------------|
| Annual | 67 | 21.90 |
| Biennial | 38 | 12.42 |
| Perennial | 201 | 65.68 |
| Total | 306 | 100.00 |

Table 7. The distribution of species in different floristic regions.

| Floristic Region | Number of Species | Percentage |
|----------------------|----------------------|------------|
| Euro-Siberian | Euro-Siberian | 68 |
| | Euxine | 14 |
| | Euxine (mt.) | 1 |
| | Hyrcano-Euxine | 3 |
| | Hyrcano-Euxine (mt.) | 1 |
| Mediterranean | Mediterranean | 26 |
| | Mediterranean (mt.) | 1 |
| | E. Mediterranean | 8 |
| Irano-Turanian | | 15 |
| More than one region | | 169 |
| Total | 306 | 100.00 |

Table 8. Comparison (%) of floristic results of the present study with those given for nearby regions in earlier studies.

| | Number of species | Euro-Siberian | Mediterranean | Irano-Turanian | More than one region | Endemic |
|-----------------------|-------------------|---------------|---------------|----------------|----------------------|---------|
| Kunduz | 306 | 28.43 | 11.44 | 4.90 | 55.23 | 4.29 |
| Özen & Kılıç (1995) | 696 | 20.75 | 13.79 | 4.32 | 61.14 | 5.18 |
| Kılıç (1985) | 850 | 23.80 | 9.50 | 16.50 | 50.20 | 2.00 |
| Kutbay & Kılıç (1995) | 611 | 25.19 | 9.65 | 2.29 | 62.87 | 3.43 |
| Karaer & Kılıç (1993) | 654 | 18.60 | 19.60 | 1.70 | 60.10 | 2.75 |
| Kılıç & Özen (1998) | 330 | 17.10 | 15.00 | 1.20 | 66.70 | - |

low due to poor ecological diversity in this area (i.e. at a given area the level of endemism is mainly correlated with ecological diversity).

A comparison of the richest families found in the present study with those in other floristic studies previously carried out near this region is given in Table 9. As seen in this table, the families *Fabaceae*, *Rosaceae*, *Asteraceae*, *Lamiaceae* and *Poaceae* are the richest in terms of taxa in the study area. In addition, these families are also placed among the top five in Turkey in previous studies carried out in the Black Sea region, Turkey.

From the geographical and phytogeographical points of view, the study area is situated in the Central Black Sea part of the Black Sea region and the Euxine sector of the Euro-Siberian floristic region. The Euxine sector has been divided into three parts from the Black Sea region to Central Anatolian as Eu-Euxine, Sub-Euxine and Xero-Euxine by Zohary (1973). According to Zohary (1973) the research area is included in the Sub-Euxine part of the Euxine sector.

Table 9. Comparison (%) of the richest families of the present study with those given for nearby regions in earlier studies.

| | Kunduz | Özen & Kilinç (1995) | Kilinç (1985) | Kutbay & Kilinç (1995) | Karaer & Kilinç (1993) | Kilinç & Özen (1998) |
|------------------|--------|----------------------|---------------|------------------------|------------------------|----------------------|
| Fabaceae | 10.78 | 9.94 | 11.30 | 10.14 | 9.63 | 12.30 |
| Rosaceae | 6.86 | 4.47 | 4.62 | 4.09 | 3.82 | 3.00 |
| Asteraceae | 6.53 | 10.38 | 11.30 | 11.62 | 10.39 | 12.30 |
| Lamiaceae | 5.88 | 5.76 | 6.25 | 6.87 | 5.19 | 7.20 |
| Poaceae | 4.90 | 7.06 | 7.97 | 6.87 | 8.25 | 6.00 |
| Brassicaceae | 3.92 | 3.89 | 5.80 | 3.60 | 3.36 | 3.60 |
| Scrophulariaceae | 3.59 | 3.46 | 4.37 | 4.41 | 3.66 | 2.40 |
| Rubiaceae | 3.26 | 2.60 | 2.50 | - | 1.37 | 1.80 |
| Boraginaceae | 2.94 | 2.74 | 2.75 | 3.10 | 3.36 | 2.40 |
| Caryophyllaceae | 2.61 | 3.31 | 4.37 | 2.61 | 3.82 | 5.10 |
| Others | 48.73 | 37.75 | 32.53 | 46.69 | 40.74 | 36.70 |

The study area includes a Mediterranaen pre-steppic intrapontic layer, a supra-Mediterranean prepontic layer, a mountainous prepontic layer, a mountainous euxine layer and a Mediterranean intrapontic layer according to the treatment by Quezel et al. (1980) using altitude and biogeography criteria.

In this study four associations are determined. One of these belongs to degraded forest vegetation and the remaining belong to forest vegetation. The phytosociological evaluation of this associations is as follows.

Carpino-Quercetum cerridis was described previously by Kutbay & Kilinç (1995), Özen & Kilinç (1995) and Quezel et al. (1980). This association is included in QUERCETEA PUBESCENTIS by Quezel et al. (1980), QUERCO-CARPINETALIA ORIENTALIS by Kutbay & Kilinç (1995) and CARPINO-ACERION by Özen & Kilinç (1995). As seen in Table 10, the associations were compared with those in similar studies by using Sorenson's similarity formula (Sorenson, 1948), as a percentage, around the Black Sea region. As in Table 10, the association *Carpino-Quercetum cerridis* resembles Özen and Kilinç's association (1995) in 48.7%.

The association *Junipero-Pinetum nigrae* is described from the Black Sea region for the first time in this paper. There are several *Pinus nigra* associations described by Akman (1972, 1973), Akman & Ketenoğlu (1978), Akman et al. (1979, 1987, 1983 a and 1983 b), Yurdakulol (1981), Akman & Aydoğdu (1986), Akman & İlarslan (1983), Özen & Kilinç (1995), Gemici (1988),

Ocakverdi (1987), Ekim & Akman (1990) and Kilinç (1985) in Turkey. This association is included in QUERCETEA PUBESCENTIS by Akman et al. (1983, 1987), Yurdakulol (1981), Akman & Aydoğdu (1986), and Akman & İlarslan (1983), in CARPINO-ACERION by Özen & Kilinç (1995), Akman et al. (1983), and Gemici (1988), in QUERCO-CEDRETTALIA LIBANI by Akman et al. (1979), Ocakverdi (1987), and Ekim & Akman (1990), and in PINO-CISTION LAURIFOLII by Kilinç (1985). The highest similarity (58.6%) to ours was obtained by Özen & Kilinç (1995) (Table 10).

The *Daphno-Pinetum sylvestris* association was named differently by some researchers (Özen & Kilinç, 1995; Kilinç, 1985; Akman et al., 1983 and 1987; Akman & Aydoğdu, 1986; Tatlı, 1987; Ekim & Akman, 1990; Düzenli, 1989). This association is included in RHODODENDRO-FAGETALIA ORIENTALIS by Özen & Kilinç (1995), in QUERCO-FAGETEA by Akman et al. (1983 and 1987), in QUERCETEA PUBESCENTIS by Akman & Aydoğdu (1986) and Düzenli (1989), in QUERCO-CEDRETTALIA LIBANI by Ekim & Akman (1990), and in CARPINO-ACERION by Kilinç (1985) and Tatlı (1987). This association is most similar to that described by Özen & Kilinç (1995) (Table 10).

The association *Galio-Fagetum orientalis* is described from the Black Sea region for the first time in this paper. There are several *Fagus orientalis* associations with different codominant species described by Özen & Kilinç (1995), Kutbay & Kilinç (1995), Quezel et al. (1980), and Akman et al. (1983) in Turkey. This association is included in FAGETALIA SYLVATICA by Akman et al.

(1983), in CARPINO-ACERION by Quezel et al. (1980), and in RHODODENDRO-FAGETALIA ORIENTALIS by Özen & Kılıç (1995) and Kutbay & Kılıç (1995). The highest similarity (49.3%) to ours was obtained by Özen & Kılıç (1995) (Table 10).

Table 10. Similarity percentages between associations in the study area and in other studies in the Black Sea region, according to Sorenson's similarity formula (Sorenson, 1948).

| | Carpino-Quercetum cerridis | Junipero-Pinetum nigrae | Daphno-Pinetum sylvestris | Galio-Fagetum orientalis |
|-----------------------|----------------------------|-------------------------|---------------------------|--------------------------|
| Özen & Kılıç (1995) | 48.70 | 58.60 | 57.10 | 49.30 |
| Kılıç (1985) | - | 16.10 | 27.30 | - |
| Kutbay & Kılıç (1995) | 30.90 | 49.30 | 38.20 | 37.50 |
| Quezel et al. (1980) | 22.80 | 23.40 | - | 23.60 |

The syntaxonomical interpretation of these associations is given below:

UPPER CLASS: *Querco-Fagea* Fukarek & Fabijanic 1968

CLASS: *Quercetea pubescens* (Oberd. 1948) doing Kraft 1955

References

- Akman Y (1972). The vegetation of Beynam Forest. *Com De la Fac Sci d'Ank Serie C Tome 16*: 29-53.
- Akman Y (1973). Apercu préliminaire sur les conditions phytocologiques de la chaîne de l'Amanos dans la région du Hatay. *Com De la Fac Sci d'Ank Serie C Tome 17*: 137-164.
- Akman Y, Ketenoğlu O (1978). The phytosociological investigations of Körögöl Mountain. *Com De la Fac Sci d'Ank Serie C2 Tome 22*: 1-24.
- Akman Y, Barbero M, Quezel P (1978). Contribution à l'étude de la végétation forestière d'Anatolie Méditerranéenne. *Phytocoenologia* 5(1): 1-79.
- Akman Y, Barbero M, Quezel P (1979). Contribution à l'étude de la végétation forestière d'Anatolie Méditerranéenne. *Phytocoenologia* 5(2-3): 189-346.
- Akman Y, Barbero M, Quezel P (1983). A phytosociological research on the vegetation of the Bolu Mountains. *Com De la Fac Sci d'Ank Serie C Tome 1*: 87-104.
- Akman Y, İlarslan R (1983). The phytosociological investigations in the district of Uluhan-Mudurnu. *Com De la Fac Sci d'Ank Serie C Tome 1*: 57-70.
- Akman Y, Yurdakulol E, Demirörs M (1983). A phytosociological research on the vegetation of the Semen Mountains (Bolu). *Com De la Fac Sci d'Ank Serie C Tome 1*: 87-104.
- ORDER: *Querco-Carpinetalia orientalis* Quezel, Barbero et Akman 1980
- ASSOCIATION: *Carpino-Quercetum cerridis* Kutbay & Kılıç 1995
- ASSOCIATION: *Junipero-Pinetum nigrae* Özen & Kılıç as. nov.
- CLASS: *Querco-Fagetea* (Br.-Bl. et Vliegler 1937) Fuk. et Fab. 1968
- ORDER: *Rhododendro-Fagetalia orientalis* Quezel, Barbero et Akman 1980
- ASSOCIATION: *Daphno-Pinetum sylvestris* Kutbay & Kılıç 1995
- ASSOCIATION: *Galio-Fagetum orientalis* Özen & Kılıç as. nov.
- Acknowledgement**
- We would like to thank Ondokuz Mayıs University Research Fund for its financial support.
- Akman Y, Aydoğdu M (1986). A phytosociological study in the vicinities of Çamlıdere, Çamkoru and Peçenek (Ankara). *Com De la Fac Sci d'Ank Serie C Tome 4*: 9-24.
- Akman Y, Quezel P, Yurdakulol E, Ketenoğlu O, Demirörs M (1987). La Vegetation des Hauts Sommets de l'Ilgaz Dağ. *Ecologia Mediterranea Tome 12 Fasc. 1/2*: 119-129.
- Akman Y (1993). *Biyolojik Anadolu*. Ankara: Palme Yayınları.
- Anonymous (1984). *Samsun ili verimlilik envanteri ve gübre ihtiyaç raporu*. Ankara: Tarım Or Köy Bak Yay No 23 Genel Yay. No 760.
- Braun-Blanquet J (1932). *Plant Sociology*. New York and London: McGraw-Hill.
- Düzenli, A (1989). Akdağmadeni (Sivas) *Pinus sylvestris* L. ormanının floristik ve fitososyolojik yapısı *Doğa Türk Bot Derg* 13 (3): 418-436.
- Davis PH (1965-1985). *Flora of Turkey and the East Aegean Islands*. Vol. 1-9. Edinburgh: Edinburgh Univ Press.
- Davis PH, Mill RR, Tan K (1988). *Flora of Turkey and the East Aegean Islands* (supplement). Vol. 10. Edinburgh: Edinburgh Univ Press.
- Donner J (1985). Verbreitungskarten zu P.H. Davis Flora of Turkey 1-8. *Linzer Biol Beitr* 17/1: 1-20.
- Donner J (1987). Verbreitungskarten zu P.H. Davis Flora of Turkey 9. *Linzer Biol Beitr* 19/1: 3-16.

- Donner J (1990). Distribution maps to P.H. Davis "Flora of Turkey" 1-10. *Linzer Biol Beitr* 22 (2): 381-515.
- Ekim T, Akman Y (1990). Eskişehir ili Sündiken Dağlarındaki orman vejetasyonunun bitki sosyolojisi bakımından araştırılması. *Doğa Tr J of Botany* 15: 28-40.
- Emberger L (1952). Surie quotienpluviometer migue. *C R Acad Sci* 2324: 2508-2510.
- Erentöz C, Kettin İ (1962) *Türkiye Jeoloji Haritası 1/500.000 ölçekli Sinop paftası*. Ankara: MTA Ens Matbaası.
- Gemicici Y (1988). Akdağ (Afyon-Denizli) ve çevresinin vejetasyonu. *Doğa Türk Botanik Derg* 12 (1): 8-57.
- Karaer F, Kilinç M (1993). Sinop Yarımadasının florası. *Doğa Tr J of Botany* 17(1): 5-20.
- Kilinç M (1985). İç Anadolu-Batı Karadeniz Geçiş Bölgesinde Devrez Çayı ile Kızılırmak Nehri Arasında Kalan Bölgenin Vejetasyonu. *Doğa Bilim Derg Seri A2* 9(2): 315-357.
- Kilinç M, Özén F (1988). Samsun Ondokuz Mayıs Üniversitesi Kurupelit kampus alanı ve çevresinin florası. *Ond May Ün Fen Derg* 1(2): 97-121.
- Kilinç M, Özén F (1993). Samsun ve Sinop'dan (A5,A6) yeni floristik kayıtlar. *Doğa Tr J of Botany* 17: 37-39.
- Kutbay HG, Kilinç M (1995). Nebyan Dağı (Samsun) ve Çevresinin Vejetasyonu Üzerinde Fitodosyolojik ve Ekolojik Bir Araştırma. *Tr J of Botany* 19(1): 41-63.
- Kutbay HG, Kilinç M (1995). Nebyan Dağı (Samsun/Bafra) florası. *Tr J of Botany* 19: 345-371.
- Ocakverdi H (1987). Seydişehir maden bölgesi (Konya) ve çevresinin vejetasyonu. *Doğa Türk Botanik Derg* 11(1): 102-148.
- Özen F, Kilinç M (1995). Alaçam-Gerze ve Boyabat-Durağan arasında kalan bölgenin florası. *Tr J of Botany* 19: 241-275.
- Özen F, Kilinç M (1995). Alaçam-Gerze ve Boyabat-Durağan arasında kalan bölgenin vejetasyonu: I-Maki, frigana, dere ve step vejetasyonları. *Tr J of Botany* 19: 65-86.
- Özen F, Kilinç M (1995). Alaçam-Gerze ve Boyabat-Durağan Arasında Kalan Bölgenin Vejetasyonu: II-Orman ve Bozuk Orman Vejetasyonları. *Tr J of Botany* 19(1): 87-105.
- Özen F (2000). New floristic records for various squares in the flora of Turkey. *BAÜ Fen Bil Ens Derg* 2: 42-46.
- Quezel P, Barbero M, Akman Y (1980). Contribution a l'étude de la vegetation forestiere d'Anatolie Septentrionale. *Phytocoenologia* 8(3/4): 365-519.
- Sorensen TA (1948). Method of establishing groups of equal amplitude in plant sociology based on similarity of species content. *Biol Skr K Danske Vidensk Selsk* 5(4): 1-34.
- Tatlı A (1987). Allahuekber Dağlarının bitki sosyolojisi yönünden araştırılması. *Doğa Türk Botanik Derg* 11(1): 169-194.
- Weber HE, Moravec J, Theurillat J (2000). Code of Phytosociological Nomenclature, 3rd edition. *Journal of Vegetation Science* 11(5): 739-767.
- Yıldırımlı Ş (1997). The chorology of Turkish species of Apiaceae family. *Ot Sist Bot Derg* 4(2): 105-128.
- Yıldırımlı Ş (1999). The chorology of Turkish species of Asteraceae family. *Ot Sist Bot Derg* 6(2): 75-123.
- Yurdakulol E (1981). A phytosociological and ecological research on the vegetation of the Pos Forests (Adana distr. Karsanti) on the Anti-Taurus Mountains. *Com De la Fac Sci d'Ank Serie C2 Tome 24*: 1-50.
- Zohary M (1973). *Geobotanical Foundations of the Middle East*. Vol. 1-2. Stuttgart: Gustav Fischer Verlag.